arXiv:1403.2557v1 [math.RA] 11 Mar 2014

Infinitely dimensional, affine nil algebras $A \otimes A^{op}$ and $A \otimes A$ exist

Agata Smoktunowicz*

March 12, 2014

Abstract

In this paper we answer two questions from [16], by showing that, over any algebraically closed field, K, there is a finitely generated, infinitely dimensional algebra A such that algebras $A \otimes_K A$ and $A \otimes_K A^{op}$ are nil.

1 Introduction

Several authors have studied the properties of tensor products of algebras over a field; for example, in [3], [2], [5], [8], [9], [11],[15] and [20]. In 1993, the question was posed in [16] as to whether every affine algebra A, such that $A \otimes A$ is nil, is locally nilpotent. In [17] it was shown that this is indeed the case for algebras over ordered fields; however, the general case remained open. In [4], a related question was asked, namely whether a Jacobson radical of algebras $A \otimes A$ and $A \otimes A^{op}$ is always locally nilpotent (recall that a nil ring is always Jacobson radical). The aim of this paper is to show that in general these two statements need not hold; in other words, that the following holds.

Theorem 1.1 *Over every algebraically closed field there is a finitely generated graded K-algebra R such that $R \otimes_K R$ and $R \otimes_K R^{op}$ are nil algebras and R is an infinitely dimensional algebra.*

The properties of algebras $A \otimes A^{op}$ may be pertinent to the study of A-bimodules. Recall that A-bimodules correspond to A^e-modules, where

*School of Mathematics, University of Edinburgh, A.Smoktunowicz@ed.ac.uk

$A^e = A \otimes A^{op}$. Determining the Jacobson radical or nil radical of algebra $A \otimes A^{op}$ may be useful in the context of the Nakayama lemma [10].

Quite often, the structure of the tensor product of an algebra with itself or with its opposite provides information about the structure of the algebra. For example, if A is a finite dimensional F-algebra such that $A \otimes A^{op}$ is a matrix F-algebra, then A is central simple. In this case, if $A \otimes A$ is a matrix F-algebra then A is a central simple of exponent 2. Several new directions have recently developed within the area of infinitely dimensional Brauer Groups. The methods developed in this paper might be related to these results, as in many cases they permit the construction of infinite dimensional graded algebras A such that $A \otimes A^{op}$ satisfy a restricted number of prescribed relations (however, in general these constructed algebras are not simple). Another application could be to construct various examples of nilpotent algebras; for some open questions in this area and connections with group theory, see [1]. Another application might be to study the properties of polynomial rings which are clean, using the results of Kanwar, Leroy and Matczuk [13], or to obtain examples of rings with clean tensor product other than those in [23].

The construction from this paper can also be applied to some other situations where we need to construct algebras A such that $A \otimes A$ and $A \otimes A^{op}$ satisfy prescribed relations; for example, it could be applied to study local rings, or rings with local tensor products $A \otimes A$ and $A \otimes A^{op}$ (for some related results, see [12]). It may also be useful for constructing examples of bialgebras or Hopf algebras satisfying some properties, for example of infinite dimensional not pointed, nil Hopf algebras; however, this seems complex.

2 Notation and plan

Let K be a field, and let A be a free algebra over K with two generators x, y of degree one. Obviously, the monomials of the form $x_{i_1} x_{i_2} \dots x_{i_n}$ where $x_{i_1}, \dots, x_{i_n} \in \{x, y\}$ form a K-basis of A. By A^1 we denote the algebra obtained from A by the adjunction of a unity. If an element $a \in A$ is a sum of monomials of the same degree multiplied by coefficients, then by $deg(a)$ we mean the number expressing the common degree of the monomials. For a positive integer n and a subset S of A by $S(n)$ we denote the set of all elements of S which are sums of monomials of length equal to n multiplied by elements from K. So $A = A(1)+A(2)+\dots$. By a slight abuse of notation, $A(0)$ will denote K (in fact it means $K \cdot 1_{A^1}$). Finally, by $\mathcal{M}$ we denote the set of all monomials of A. We put $A(\alpha) = 0$ if $\alpha < 0$. An ideal in A will

mean a two-sided ideal in A. Let S be a linear space over a field K, then $\dim_K S$ will denote the dimension of S. We say that element $a \in A \otimes A$ is homogeneous if $a \in A(m) \otimes A(n)$ for some m, n. We will then say that the left degree of a is m, denoted $l(a) = m$, and the right degree of a is n, denoted as $r(a) = n$. A linear subspace of $A \otimes A$ is homogeneous if it is generated by homogeneous elements.

We will use the following notation throughout this paper.

Let $1 \leq \alpha$, $0 < n_1 < n_2$ be natural numbers and let $F \subseteq A(n_1) \otimes A(\alpha)$, $G \subseteq A(\alpha) \otimes A(n_1)$. We define

$$Q(F, n_1, n_2) = \sum_{k=0}^{\infty} \sum_{j \in W(n_1,n_2)} ((A(k \cdot n_2 - n_1) + A(k \cdot n_2)) \otimes A(j))F(A \otimes A),$$

$$Q'(G, n_1, n_2) = \sum_{k=0}^{\infty} \sum_{j \in W(n_1,n_2)} (A(j) \otimes (A(k \cdot n_2) + A(k \cdot n_2 - n_1))G(A \otimes A),$$

where $j \in W(n_1, n_2)$ if and only if the interval $[j, j + \alpha]$ is disjoint with all intervals $[kn_2 - n_1, kn_2 + n_1]$ for $k = 0, 1, 2, \ldots$.

An outline of the proof now follows.

- In Theorem 3.1 we will find a nil ideal N in $A \otimes A$ and appropriate linear subspaces $S_i \subseteq A \otimes A$ and such that $N \subseteq \sum_{i=1}^{\infty} S_i$. The proof of Theorem 3.1 will be presented in sections 9-12 in the Appendix at the end of this paper.

- In Theorem 6.1 we find linear spaces $U_i \subseteq A(a_i)$ such that roughly speaking $S_i \subseteq A \otimes U_i + U_i \otimes A$ for $i = 1, 2, \ldots$. In Section 4 linear mappings are introduced, which are important for the construction of sets U_i.

- In Sections 5 and 6 the Golod-Shafarevich Construction and Hilbert Nullstellensatz Theorem are used, to assure that the codimesion of U_i in $A(a_i)$ is 1.

- In Corollary 6.1, using sets U_i, special linear space D will be constructed such that $N \subseteq D \otimes A + A \otimes D$.

- In Section 7 we define $I[D] = \{r \in D : Ar \subseteq D\}$. Here it is shown that $I[D]$ is an ideal in A.

- In Section 11.3 it is shown that $A/I[D] \otimes (A/I[D])^{op}$ and $A/I[D] \otimes (A/I[D])$ are nil algebras.

3 Embedding nil ideals in linear spaces

The aim of this section is to demonstrate that some nil ideals in $A \otimes A$ are contained in the sum of appropriate linear spaces S_i.

Let $x_i \in \{x, y\}$ be generators of A, then as usual we write $(x_1 x_2 \cdots x_n)^{op} = x_n x_{n-1} \cdots x_1$. Let $a_i, b_i \in A$. For $r = \sum_i a_i \otimes b_i$ define $\xi(r) = \sum_i a_i \otimes (b_i)^{op}$.

Theorem 3.1 *Let K be a field and let A be a K-algebra generated in degree one by two elements. Let $0 < a_1 < a_2 < \ldots$ be integers such that a_i divides a_{i+1}, $a_{i+1} > 100ia_i$, for every $i = 1, 2, \ldots$, and $a_1 > 100$. Then, for all $i \geq 1$ there exist homogeneous linear spaces $G_i \subseteq A(a_i) \otimes A(a_i)$, $F_i \subseteq A(a_i) \otimes \sum_{j > \frac{a_i}{10i}}^{10ia_i} A(j)$, $F_i' \subseteq \sum_{j > \frac{a_i}{10i}}^{10ia_i} A(j) \otimes A(a_i)$ with $\dim_K G_i, \dim_K F_i, \dim_K F_i' < 2^{500ia_{i-1}} a_{i+2}^{100i}$, also $G_1 = 0, G_2 = 0$, $F_1 = 0$, $F_1' = 0$. Moreover, there exist an ideal N in $A \otimes A$ such that $N \subseteq \sum_{n=1}^{\infty} S_n$ where S_n is a linear space such that $S_n = \sum_{n=1}^{\infty} M_n + Q_n + Q_{n-1}$ and where*

$$M_n = \sum_{0 \leq k, k'} (A(ka_{n+1}) \otimes (A(k'a_{n+1}) + A(k'a_{n+1} - a_n)))G_n(A \otimes A)$$

$$Q_n = Q(F_n, a_n, a_{n+1}) + Q'(F_n', a_n, a_{n+1}), Q_0 = 0.$$

Moreover, for every $f \in A \otimes A$ there is $n = n(f)$ such that $f^n, \xi(f^n) \in N$.

Proof. The proof is long and technical, and in most cases follows ideas from [21]. The detailed proof is presented in the Appendix, sections 9-12 at the end of this paper. Observe that ξ is defined at the beginning of this theorem. □

Notice that f_i in Theorem 3.1 need not be nonzero.

4 Linear mappings

In this section we will introduce sets $U_i \subseteq A(a_i)$ and sets $B_{a_{i+1}}(U_i)$, which will be related to sets Q_1, Q_2, Q_3 from Theorem 3.1. Throughout this paper we will use the following notation. Let n, m be natural numbers and $m < n$. Given set $U \subseteq A(m)$ define

$$B_n(U) = \sum_{k=0}^{\infty} A(k \cdot n) U_i A + A(k \cdot n - m) U_i A.$$

For each i, denote

$$A[a_{i+1}] = \sum_{k=1}^{\infty} A(k \cdot a_{i+1}).$$

Given linear mapping f, by $kerf$ we will denote the kernel of f. We begin with the following Lemma.

Lemma 4.1 *Let $a_1, a_2, \ldots$ be a sequence of natural numbers such that $2a_i < a_{i+1}$ and a_i divides a_{i+1} for all i, and $a_0 = 0$. Suppose that for $1 \leq i \leq n$ we are given a linear space $U_i \subseteq A(a_i)$, and a monomial $m_i \in A(a_i)$ such that*

$$U_i \oplus Km_i = A(a_i), A(a_i) \cap \sum_{j=1}^{i-1} B_{a_{j+1}}(U_j) \subseteq U_i.$$

Then for $2 \leq i \leq n+1$ there are linear mappings $\delta_i : A[a_i] \to A$ such that for all $k \geq 0, i \geq 2$,

$$\delta_i(A(k \cdot a_i)) = A(k \cdot a_i \cdot t_i), A[a_i] \cap \sum_{j=1}^{i-1} B_{a_{j+1}}(U_i) \subseteq ker\delta_i$$

where $t_i = \prod_{j=1}^{i-1}(1 - \frac{2a_j}{a_{j+1}})$.

Proof. Induction by n. For $n = 2$ we define δ_2 in the following way. Let $\gamma_1 : A(a_1) \to K$ be a linear mapping with kernel equal to U_1 and such that $\gamma_1(m_1) = 1$. Let $\delta_2 : A(a_2) \to A$ be a linear mapping defined as follows, if $v = v_1v_2v_3$ for monomials $v_1, v_3 \in M(a_1)$ and $v_2 \in M(a_2 - 2a_1)$, then $\delta_2(v) = \gamma_1(v_1) \cdot v_2 \cdot \gamma_1(v_3)$. We then extend δ_2 by linearity to all elements of $A(a_2)$.

We now extend mapping δ_2 by linearity to all homogeneous elements with degrees divisible by a_2. Let $w = \prod_{i=1}^{k} w_i$ with all $w_i \in M(a_2)$. We can now define $\delta_2(w) = \prod_{i=1}^{k} \delta_2(w_i)$, and extend δ_2 by linearity to all elements from $A[a_2]$.

Observe that $\delta_2(A(a_2)) = A(a_2 - 2a_1) = A(a_2(1 - \frac{2a_1}{a_2}))$, so $\delta_2(A(k \cdot a_2)) = A(k \cdot a_2 t_2)$, as required.

Observe that if $p \in B_{a_2}(U_1)$ and p is a homogeneous element of degree divisible by a_2, then $\delta_2(p) = 0$. It follows because every $p \in B_{a_2}(U_1)$ is a linear combination of elements from $M(ka_2)U_1M$ and from $M(ka_2)M(a_2 - a_1)U_1M$, images of such elements under δ_2 are zero.

Suppose now that for every $i \leq n$ we defined mappings $\delta_i : A[a_i] \to A[a_i]$ in a such way that for every $i \leq n$ we have

$$\delta_i(A(k \cdot a_i)) = A(k \cdot a_i \cdot t_i), A[a_i] \cap \sum_{j=1}^{i-1} B_{a_{j+1}}(U_i) \subseteq ker\delta_i.$$

To construct δ_{n+1} we first define $\gamma_n : A(a_n) \to A(a_n)$. Let γ_n be a linear mapping which has kernel U_n and $\gamma_n(m_{n+1}) = 1$. Let $\delta_{n+1} : A(a_{n+1}) \to$

$A(a_{n+1})$ be a linear map defined as follows, if $v = v_1v_2v_3$ for monomials $v_1, v_3 \in M(a_n)$ and $v_2 \in M(a_{n+1} - 2a_n)$, then $\delta_{n+1}(v) = \gamma_n(v_1)\delta_n(v_2)\gamma_n(v_3)$. We then extend δ_{n+1} by linearity to all elements of $A(a_{n+1})$.

We now extend mapping δ_{n+1} to elements with degrees divisible by a_{n+1} in the following way. Let $w = \prod_{i=1}^{k} w_i$ with all $w_i \in A(a_{n+1})$. We can now define $\delta_{n+1}(w) = \prod_{i=1}^{k} \delta_{n+1}(w_i)$, and extend δ_{n+1} by linearity to all elements from $A[a_n]$.

Observe that $\delta_{n+1}(A(a_{n+1})) = \delta_n(A(a_{n+1} - 2a_n)) = A((a_{n+1} - 2a_n)t_n) = A(a_{n+1}(1 - \frac{2a_n}{a_{n+1}})t_n) = A(a_{n+1}t_{n+1})$, as required.

We will now show that if $p \in \sum_{i=1}^{n} B_{a_{i+1}}(U_i)$ and p is a homogeneous element of degree divisible by a_{n+1}, then $\delta_{n+1}(p) = 0$. By the inductive assumption we may assume that if $p' \in \sum_{i=1}^{n-1} B(U_i)$ and p' is a homogeneous element of degree divisible by a_n, then $\delta_n(p') = 0$. As δ_{n+1} is a linear mapping, it suffices to consider the following two cases only.

Case 1. $p \in B$ where $B = \sum_{i=1}^{n-1} B_{a_{i+1}}(U_i)$, If $p \in A(k \cdot a_{n+1})$ and $p \in B$, then p is a linear combination of elements from $M(ka_{n+1})rM$ where $r \in A(a_{n+1})$ and either $r \in (A(a_n) \cap B)M(a_{n+1} - a_n)$, or $r \in M(a_n)(A(a_{n+1} - 2a_n) \cap B)M(a_n)$ or $r \in M(a_{n+1} - a_n)(A(a_n) \cap B)$ (because a_n divides a_{n+1}). By Assumptions $B \cap A(a_n) \subseteq U_n \subseteq ker\gamma_n$. By the definition of the mapping δ_{n+1}, we get $\delta_{n+1}(r) = 0$ in all these cases, because $\delta_n(B \cap A(k \cdot a_n)) = 0$ by the inductive assumption.

Case 2. $p \in B_{a_{n+1}}(U_n)$. Observe now that if $p \in A(k \cdot a_{n+1})$ and $p \in B_{a_{n+1}}(U_n)$ then p is a linear combination of elements from $M(ka_{n+1})rM$ where $r \in A(a_{n+1})$ and either $r \in U_nM(a_{n+1} - a_n)$, or $r \in A(a_{n+1} - a_n)U_n$. Recall that $\gamma_n(U_n) = 0$. By the definition of the mapping δ_{n+1} we get $\delta_n(r) = 0$ in all these cases. □

Let m, n, α be natural numbers with $m > n$, and let $g \in A(\alpha)$. Define

$$M(g, n, m) = \sum_{i:[i,i+\alpha] \in W} A(i)gA$$

where $[i, i + \alpha] \in W$ if and only if the interval $[i, i + \alpha]$ is disjoint with interval $[km - n, km + n]$, for every $k \geq 0$. In Section 6 we will use the fact that $M(g, n, m)$ are related to some linear spaces $Q(f, n, m)$, $Q'(f, n, m)$.

Lemma 4.2 *Let $a_1, a_2, \ldots$ be a sequence of natural numbers such that $0 < 2000j^2a_j < a_{j+1}$ and $10(j+1)^2a_j$ divides a_{j+1} for all $j \geq 1$, and 10 divides a_1. Let i be a natural number. Suppose that for all $1 \leq j \leq i$ we are given linear spaces $U_j \subseteq A(a_j)$ and monomials $m_j \in A(a_j)$ such that $U_j \oplus Km_j = A(a_j)$ and $A(a_j) \cap \sum_{k=1}^{j-1} B_{a_{k+1}}(U_k) \subseteq U_j$.*

Let $n \leq i$ and let $f \in A(\alpha)$ for some $\frac{a_n}{10n} \leq \alpha \leq 10n \cdot a_n$. Denote $F = M(f, a_n, a_{n+1})$. Suppose that $F \cap A(a_{n+1+k}) \subseteq U_{n+1+k}$, for all $k \geq 0$,

with $n+k+1 \leq i$. Let mappings δ_j be constructed as in the proof of Lemma 4.1. Then there is a set $E(f)$ with cardinality smaller than $2a_n^4 \cdot 2^{2a_{n-1}}$, and such that $\delta_{n+1+k}(F \cap A[a_{n+1+k}]) \subseteq < E(f) >$, for all $k \geq 0$ with $k+n+1 \leq i+1$ where $< E(f) >$ is the ideal generated in A by elements from $E(f)$. Moreover, $E(f)$ is a homogeneous set and $E(f) \subseteq \sum_{i > \frac{a_n}{40n}}^{20na_n} A(i)$.

Proof. Recall that $f \in A(\alpha)$. Denote first $H(f) = \{\delta_n(M(i)fM(j)) : i, j < a_n$, and a_n divides $i + \alpha + j\}$. Then $H(f) \subseteq \delta_n(\sum_{i=\frac{a_n}{10n}}^{10na_n} A(i)) \subseteq \sum_{i=\frac{t_n a_n}{10n}}^{10nt_n a_n} A(i)$. We will first show that $\delta_{n+1+k}(F \cap A[a_{n+1+k}]) \subseteq < H(f) >$, for all $k \geq 0$ with $k+n+1 \leq i+1$ where $< H(f) >$ is the ideal generated in A by elements from $H(f)$.

We will first show that $\delta_{n+1}(F \cap A[a_{n+1}]) \subseteq < H(f) >$. Recall that elements from $F \cap A[a_{n+1}]$ consist of linear combination of elements from $M(j)fM(j')$ where $j + \alpha + j' = \beta \cdot a_{n+1}$, for some β and the interval $[j, j+\alpha]$ is disjoint with any interval $[ka_{n+1} - a_n, ka_{n+1} + a_n.]$ We can write $j = c + c'$, $j' = d + d'$ where c', d' are divisible by a_{n+1} and $0 \leq c, d < a_{n+1}$. As $\alpha \leq \frac{1}{200} a_{n+1}$ and $c + \alpha + d$ is divisible by a_{n+1} it follows that either $c+\alpha+d = a_{n+1}$ or $c+\alpha+d = 2a_{n+1}$. If the later holds then $c, d \geq a_{n+1} - \alpha$ which implies that the interval $[c, c+\alpha]$ is not disjoint with interval $[a_{n+1} - a_n, a_{n+1} + a_n]$, hence the interval $[j, j+\alpha]$ is not disjoint with some interval $[ka_{n+1} - a_n, ka_{n+1} + a_n.]$ It follows that $c+\alpha+d = a_{n+1}$. Observe now that, by the definition of δ_{n+1}, and by Lemma 4.1 we get $\delta_{n+1}(M(j)fM(j')) = \delta_{n+1}(M(c'))\delta_{n+1}(M(c)fM(d))\delta_{n+1}(M(d')) \subseteq A\delta_{n+1}(M(c)fM(d))A$. Consequently in is sufficient to show that $\delta_{n+1}(M(c)fM(d)) \subseteq < H(f) >$.

Recall that $c + \alpha + d = a_{n+1}$ and the interval $[c, c+\alpha]$ is disjoint with interval $[a_{n+1} - a_n, a_{n+1} + a_n]$, so $\deg d > a_n$. Observe also that $\deg c > a_n$ because the interval $[c, c+\alpha]$ is disjoint with interval $[-a_n, a_n]$. We can write $c = c_1 c_2$, $d = d_1 d_2$ where $c_1, d_2 \in M(a_n)$. By the definition of σ_{n+1} we get $\delta_{n+1}(cfd) = \delta_{n+1}(c_1 c_2 f d_1 d_2) = \gamma_n(c_1)\delta_n(c_2 f d_1)\gamma_n(d_2) \in K \cdot \delta_n(c_2 f d_1)$.

We can write $c_2 = c_3 + c_4$, $d_1 = d_3 + d_4$ where c_3, d_4 are divisible by a_n, and c_4, d_3 are monomials of degres less than a_n. By the definition of δ_n we then have $\delta_n(M(c_2)fM(d_1)) = \delta_n(M(c_3))\delta_n(M(c_4)fM(d_3))\delta(M(d_4)) \subseteq A(c_3 t_n)\delta_n(M(c_4)fM(d_3))A(d_4 t_n)$. By the definition of $H(f)$ at the beginning of this proof $\delta_n(M(c_4)fM(d_3)) \in H(f)$, as required.

We will now show that $\delta_{n+1+k}(F \cap A[a_{n+1+k}]) \subseteq < H(f) >$, for all $k \geq 0$, by induction on k. If $k = 0$ the result follows from the first part of this proof. Suppose that $k \geq 1$ and that it is true for all numbers smaller than k. By the inductive assumption $\delta_{n+k}(F \cap A[a_{n+k}]) \subseteq < H(f) >$. We need to show that $\delta_{n+k+1}(F \cap A[a_{n+k+1}]) \subseteq < H(f) >$. Recall that elements from $F \cap A[a_{n+1}]$ consist of linear combination of elements from $M(j)fM(j')$

where $j+\alpha+j' = \beta \cdot a_{n+k+1}$, for some β and the interval $[j, j+\alpha]$ is disjoint with any interval $[la_{n+1}-a_n, la_{n+1}+a_n]$ (and hence disjoint with any interval $[la_{n+k}-a_n, la_{n+k}+a_n]$, and any interval $[la_{n+k+1}-a_n, la_{n+k+1}+a_n]$.)

Similarly as before it suffices to consider the case when $j, j' < a_{n+k+1}$ and either $j+j'+\alpha = a_{n+k+1}$ or $j+j'+\alpha = 2a_{n+k+1}$. If the later holds then, $j, j' \geq a_{n+k+1}-\alpha$. Therefore $[j, j+\alpha]$ contains point a_{n+k+1} and $j < a_{n+k+1}$. Therefore, the interval $[j, j+\alpha]$ is not disjoint with interval $[a_{n+k+1}-a_n, a_{n+k+1}+a_n]$, a contradiction. It follows that $j+j'+\alpha = a_{n+k+1}$. Observe that interval $[j, j+\alpha]$ cannot contain any point $l \cdot a_{n+k}$, because then it would not be disjoint with some interval $[l \cdot a_{n+k}-a_n, l \cdot a_{n+k}+a_n]$. It follows that either $j+\alpha < a_{n+k}$; or $j > a_{n+k}$ and $j+\alpha < a_{n+k+1}-a_{n+k}$; or $j > a_{n+k+1}-a_{n+k}$.

By the definition of δ_{n+k+1}, if $j+\alpha < a_{n+k}$ then $\delta_{n+k+1}(M(j)fM(j')) = \gamma_{n+k}(M(j)fM(a_{n+k}-j-\alpha))\delta_{n+k}(M(a_{n+k+1}-2a_{n+k}))\gamma_{n+k}(M(a_{n+k}))$. As $M(j)fM(a_{n+k}-j-\alpha) \subseteq F \cap A(a_{n+k}) \subseteq U_{n+k} \subseteq ker\gamma_{n+k}$ by assumption, so $\delta_{n+k+1}(M(j)fM(j')) = 0$.

If $j > a_{n+k}$ and $j+\alpha < a_{n+k+1}-a_{n+k}$ then $\delta_{n+k+1}(M(j)fM(j')) = \gamma_{n+k}(M(a_{n+k}))\delta_{n+k}(M(j-a_{n+k})fM(j'-a_{n+k}))\gamma_{n+k}(M(a_{n+k}))$. By the inductive assumption $\delta_{n+k}(M(j-a_{n+k})fM(j'-a_{n+k})) \subseteq < H(f) >$, hence we get $\delta_{n+k+1}(M(j)fM(j')) \subseteq < H(f) >$.

If $j > a_{n+k+1}-a_{n+k}$ then $\alpha+j' < a_{n+k}$. Hence, $\delta_{n+k+1}(M(j)fM(j')) = \gamma_{n+k}(M(a_{n+k}))\delta_{n+k}(M(a_{n+k+1}-2a_{n+k}))\gamma_{n+k}(M(j-a_{n+k+1}+a_{n+k})fM(j')) = 0$, since $M(j-a_{n+k+1}+a_{n+k})fM(j') \subseteq F \cap A(a_{n+k}) \subseteq U_{n+k} \subseteq \ker\gamma_{n+k}$.

We have shown that $\delta_{n+1+k}(F \cap A[a_{n+1+k}]) \subseteq < H(f) >$, for all $k \geq 0$ with $k+n+1 \leq i+1$.

Assume that c is a natural number divisible by a_{n-1}, and that c is smaller than a_n. Define $h_c : A(c) \to A$ be a function such that: if $c = a_{n-1}$ then $h_c = \gamma_{n-1}$; if $a_{n-1} < c < a_n$ then for a monomial $v = v_1v_2$ with $v_1 \in M(c-a_{n-1})$, $v_2 \in M(a_{n-1})$ we define $h_c(v_1v_2) = \delta_{n-1}(v_1)\gamma_{n-1}(v_2)$, we then extend h_c by linearity to all elements of $A(c)$ (where γ_{n-1} is as in the definition of the mapping δ_n). Similarly, define for such c, function $h'_c : A(c) \to A$ in the following way: if $c = a_{n-1}$ then $h_c = \gamma_{n-1}$; if $a_{n-1} < c < a_n$ then for a monomial $v = v_2v_1$ with $v_1 \in M(c-a_{n-1})$, $v_2 \in M(a_{n-1})$ we define $h_c(v_1v_2) = \gamma_{n-1}(v_2)\delta_{n-1}(v_1)$, we then extend h_c by linearity to all elements of $A(c)$. We define $h(1) = h'(1) = 1$ if $r = 1$.

Let now p be a natural number divisible by a_n and let $0 \leq c, d < a_n$, $p-c-d$ is divisible by a_n (we will set $h_{p,c,d} = 0$ if $p-c-d$ is not divisible by a_n). Define function $g_{p,c,d} : A(p) \to A$ as follows first for monomials: if $v_1 \in M(c), v_2 \in M(p-c-d), v_3 \in A(d)$ we define $h_{p,c,d} = h_c(v_1)\delta_n(v_2)h'_d(v_3)$, we then extend $h_{p,c,d}$ by linearity to all elements from $A(p)$.

Let $X = \{M(i)fM(j) : 0 \leq i,j < a_{n-1}$, and a_{n-1} divides $i+\alpha+j\}$. Observe that $X \subseteq \sum_{i=\alpha}^{\alpha+2a_{n-1}-2} A(i)$, and that the cardinality of X is less than $2^{2a_{n-1}}$. Define

$$E(f) = \{h_{p,c,d}(r) : r \in X, p \in [\alpha, \alpha+2a_{n-1}-2], 0 \leq c,d < a_{n-1}\}$$

Observe that the set $H(f)$ from the beginning of our proof is spanned by elements of the form vfv' where v, v' are monomials of degree less than a_n. Write $v = wu$, $v' = u'w'$ where u, u' are monomials of degree $< a_{n-1}$ and w, w' are monomials of degrees t, t' with t, t' divisible by a_{n-1}. Then by the definition of function δ_n we get $\delta_n(vfv') \in h'_t(w) \cdot h_{p,c,d}(ufu')h_{t'}(w')$ where $c = a_n - t$ if $t > 0$ and $c = 0$ if $t = 0$; and $c' = a_n - t'$ if $t' > 0$ and $c' = 0$ if $t' = 0$ (because vfv' has degree divisible by a_n since $vfv' \in H(f)$). It follows that $H(f) \subseteq < E(f) >$.

Observe that the cardinality of $E(f)$ depends on the number of choices of numbers c, d of degrees less than a_n and on the number of choices of p, and of the cardinality of X. Since $p \in [\alpha, \alpha+2a_{n-1}-2]$ we get that the cardinality of $E(f)$ is smaller than $a_n^2 \cdot 2a_{n-1} \cdot 2^{2a_{n-1}} < 2a_n^4 2^{2a_{n-1}}$.

Recall that $\delta_n(A(a_n)) = A(t_n \cdot a_n)$. Observe that if $p - c - d \geq 0$ then $p - c - d \geq a_n$ and so $h_{p,c,d}(r) \in \sum_{i=a_n t_n}^{\alpha+2a_{n-1}} A(i)$, for all $r \in X$. Recall that $\delta_n(A(k \cdot a_{n-1}) = A(k \cdot t_{n-1} \cdot a_{n-1})$. If $p - c - d = 0$ then $h_{p,c,d}(r) \in \sum_{i=(\alpha-2a_{n-1})t_{n-1}}^{\alpha+2a_{n-1}} A(i)$, for all $r \in X$. Recall that senence $a_1, a_2, \ldots$ grows very guickly. It can be inductively proved that $t_n > \frac{1}{2} + \frac{1}{3n}$. Recall that $\alpha > \frac{a_n}{10n}$. It follows that $E(f)$ is a homogeneous set and $E(f) \subseteq \sum_{i > \frac{a_n}{40n}}^{20na_n} A(i)$. □

5 A short section on the Golod-Shafarevich theorem.

We begin by recalling Theorem 2.1 from [18] for the special case when $d = 2$. In our situation $A = T = K[x_1, x_2]$, we use algebra A with two generators x, y, so $d = 2$.

Theorem 5.1 *[Theorem 2.1, [18]] Let $I \subseteq A$ be a homogeneous two-sided ideal $I = \oplus_{n=1,2,\ldots} I(n)$. Write $A(n) = I(n) \oplus B(n)$ where $b_n = \dim_K B(n)$. Suppose that ideal I is generated by homogeneous elements $f_1, f_2, \ldots \in A$. Let r_l be the number of f_j's of degree l (thus $r_1 = 0$). Then for all $n \geq 2$*

$$b_n \geq db_{n-1} - \sum_{j=0}^{n-2} r_{n-j} b_j.$$

As mentioned in [18], at the heart of the Golod-Shafarevich construction is the following proposition.

Proposition 5.1 *[6] Let b_n, r_l be the sequences in Theorem 5.1. Let $e > 0$ be such that $e < \frac{1}{2}$. If $r_l \leq e^2(2-2e)^{l-2}$ for all $l \geq 2$, then A/I is an infinitely dimensional algebra.*

A simple proof of this proposition can be found in [18], proof of Proposition 3.1.

We will now apply these lemmas to prove Lemma 5.1.

Recall that $t_i = \prod_{j=1}^{i-1}(1 - \frac{2a_j}{a_{j+1}})$.

Lemma 5.1 *There are natural numbers $0 < a_1 < a_2 < \ldots$ such that $200(i+1)^2 a_i$ divides a_{i+1} for every $i \geq 1$, which satisfy the following condition: If I is an ideal generated by $2^{510na_{n-1}}a_{n+2}^{110n}$ with degrees from $\sum_{i>\frac{a_n}{40n}}^{20na_n} A(i)$ then A/I is an infinitely dimensional algebra.*

Proof. It can be inductively proved that $t_i > \frac{1}{2} + \frac{1}{3i}$, for all $i > 1$. Let $a_1 > 1000000^{1000^{1000}}$ and define inductively

$$a_{i+1} = a_i \cdot (1000(i+1))^{110}.$$

Let r_l denote the number of homogeneous generating relations of degree l in I. So if $\frac{a_n}{40n} \leq l \leq 20na_n$ then $r_l < 2^{510na_{n-1}}a_{n+2}^{110n}$ Denote $e = \frac{2-2^{\frac{1}{4}}}{2}$, then $2 - 2e = 2^{\frac{1}{4}}$. Denote $\xi = 2^{\frac{1}{8}}$, so

$$\xi^2 = e.$$

Consequently, to show that $r_l < e^2(2-2e)^l$ for every l, it suffices to show that for every $n \geq 1$

$$2^{510na_{n-1}}a_{n+2}^{110n} < e^2(2-e)^{\frac{a_n}{40n}-2}.$$

We will first show that $2^{510na_{n-1}} < e^2\xi^{\frac{a_n}{40n}-4}$. Recall that $a_n = a_{n-1}(1000n)^{100}$, so $\xi^{\frac{a_n}{40n}} = 2^{\frac{1}{8}\cdot\frac{1}{40n}\cdot(a_{n-1}\cdot(1000n)^{100})}$. Notice that $e^2 > \xi^{-80}$. It follows that $e^2\xi^{\frac{a_n}{40n}-4} > 2^{510na_{n-1}}$.

We will now show that $a_{n+2}^{110n} < \xi^{\frac{a_n}{40n}}$. Observe first that $\xi^{\frac{a_n}{40n}} = 2^{\frac{a_n}{320n}}$ and $a_{n+2} = a_n \cdot (1000(n+1))^{100} \cdot (1000(n+2))^{100}$. Observe that $a_{n+2} < a_n^2$, for $n = 1, 2, 3, 4, 5, 6, 7$ it follows since a_1 is very large, and for bigger n it follows since $a_n = a_1 \cdot (\prod_{i=2}^{n}(1000i))^{1000} > a_1 \cdot (1000n)^{100\cdot\frac{n}{2}-3}$. So it suffices to show that ${a_n}^{220n} < 2^{\frac{a_n}{320n}}$, or equivalently that $\sqrt{a_n} < 2^{\frac{a_n}{320n}\cdot\frac{1}{440n}}$, Observe now that $\frac{a_n}{320n} \cdot \frac{1}{440n} > \sqrt{a_n}$. Therefore it suffices to show that $c < 2^c$ where $c = \sqrt{a_n}$, and ithis is true for all $c \geq 1$.

Hence we proved that $a_{n+2}^{110n} < \xi^{\frac{a_n}{40n}}$.

As $\xi^2 = 2 - e$ it follows that $2^{510na_{n-1}} a_{n+2}^{110n} < e^2 \xi^{\frac{a_n}{40n}-4} \cdot \xi^{\frac{a_n}{40n}} = e^2(2-e)^{\frac{a_n}{40n}-2}$ for every $n \geq 1$. It follows that I is generated by $r_m < e^2 \xi^{m-2}$ homogeneous relations of degree m, for each $m > 1$. By Proposition 5.1, A/I is an infinitely dimensional algebra. □

We will now introduce a simple lemma.

Lemma 5.2 *Suppose that r_n satisfy assumptions of Proposition 5.1 for each $i > 1$. Let J be an arbitrary ideal generated by s_n homogeneous elements of degree n, with $s_n \leq r_n$ for each $n > 1$. Then for every m*

$$dim_K A(m)/J(m) > r_m - s_m.$$

Proof. Observe first that by Proposition 5.1, A/I is an infinitely dimensional algebra, for every ideal I generated by r_i homogeneous elements of degree i, for every $i > 1$.

Suppose on the contrary that $\dim_K A(m)/J(m) \leq r_m - s_m$. Observe that there is ideal I generated by at most r_i relations of degree i for each $i \geq 2$ and such that $J \subseteq I$. Moreover, in the degree m ideal I is generated by s_m elements from J and by $r_m - s_m$ elements from $A(m)$ which correspond to the basis of $A(m)/J(m)$ (so they are equal to elements of this basis modulo $J(m)$). Then $A(m)/I(m) = 0$, and so A/I is nilpotent, a contradiction with assumption that A/I is infinitely dimensional. □

Lemma 5.3 *There exist natural numbers $0 < a_1 < a_2 < a_3 < \ldots$ such that $200(i+1)^2 a_i$ divides a_{i+1} for every $i \geq 1$, which satisfy the following condition: if J is an arbitrary ideal in A generated by elements from arbitrary sets $E(Z_n) \subseteq \sum_{i=\frac{a_n}{40n}}^{20na_n} A(i)$, with cardinality of $E(Z_n)$ less than $2^{505na_{n-1}} a_{n+2}^{110n}$ for all $n \geq 1$, then*

$$\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap J) > 4 \cdot (2^{500(i+1)a_i} a_{i+3}^{100(i+1)}) + 2.$$

In particular, if G_{i+1} is as in Lemma 6.1 then

$$\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap J) > 4 \cdot \dim_K G_{i+1} + 2.$$

Proof. Define J to be an ideal generated by elements from sets $E(Z_n)$ for all n. Let s_m be the number of generating relations of degree m in J. Let I be an ideal generated by relations from sets $E(Z_n)$ and also by $4 \cdot (2^{500na_{n-1}} a_{n+2}^{100n}) + 2$ homogeneous relations of degree $t_n a_n$ for each $n \geq 1$.

Notice that $(4 \cdot (2^{500na_{n-1}} a_{n+2}^{100n}) + 2) + 2^{505na_{n-1}} a_{n+2}^{110n} < 2^{510na_{n-1}} a_{n+2}^{110n}$. Therefore I satisfies assumptions of Lemma 5.1 (since $\frac{1}{2} < t_n \leq 1$). By

Lemma 5.1 A/I is infinitely dimensional algebra. Moreover, by the proof of Lemma 5.1, I is generated by r_m homogeneous relations of degree m for all $m \geq 0$, with r_m satisfying assumptions of Proposition 5.1.

By Lemma 5.2 for this I and J we have $dim_K A(m)/J(m) > r_m - s_m$. For $m = t_{i+1}a_{i+1}$ we get that $\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap J > r_{t_{i+1}a_{i+1}} - s_{t_{i+1}a_{i+1}}$.

By construction of I, $r_{t_{n+1}a_{i+1}} - s_{t_{n+1}a_{i+1}} = 4 \cdot (2^{500(i+1)a_i} a_{i+3}^{100(i+1)}) + 2$. Therefore, $\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap J) > 4 \cdot (2^{500(i+1)a_i} a_{i+3}^{100(i+1)}) + 2$. Since $dim_K G_{i+1} \leq 2^{500(i+1)a_i} a_{i+3}^{100(i+1)}$, we get $\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap J) > 4 \cdot \dim_K G_{i+1} + 2$. □

6 Applying Golod-Shafarevich theorem and Hilbert Nullstellensatz theorem.

In this section we first give supporting lemmas, and then prove Theorem 6.1, which will, roughly speaking, show that linear spaces S_i from Theorem 3.1 are contained in $U_i \otimes_K A + A \otimes_K U_i$, for suitable linear spaces $U_i \subseteq A$.

We begin with the following lemma, which closely resembles Lemma 4.3 [22] and has the same proof.

Lemma 6.1 *Let K be an algebraically closed field, let m be a natural number, and let $T \subseteq A(m)$ and $Q \subseteq A(m) \otimes_K A(m)$ be K-linear spaces such that $\dim_K T + 4\dim_K Q \leq \dim_K A(m) - 2$.*

Then there exists a linear space $F \subseteq A(m)$ such that $T \subseteq F$ and $Q \subseteq F \otimes_K A(m) + A(m) \otimes_K F$, and $\dim_K A(m) = 1 + \dim_K U$.

Proof. The same proof as in Lemma 4.3 in [22] can be used. Notice that the fact that $m = 2^n$ in Lemma 4.3 [22] is not used in the proof, so the proof works for all m. Also in the proof variable Z can be put to be zero to make it easier. The main idea of the proof is to use the Hilbert Nullstellensatz theorem. □

Lemma 6.2 *Let i be a natural number, and $a_{i+1} < a_{i+2}$ be natural numbers such that $200(i+1)^2 a_{i+1}$ divides a_{i+2}. Let $U_{i+1}, F_{i+1}, F'_{i+1}$ be linear spaces, and let $U_{i+1} \subseteq A(a_{i+1})$, $m_{i+1} \in M(a_{i+1})$ be such that $U_{i+1} + Km_{i+1} = A(a_{i+1})$. Moreover, let $F_{i+1} \subseteq A(a_{i+1}) \otimes \sum_{j=\frac{a_{i+1}}{10(i+1)}}^{10(i+1)a_{i+1}} A(j)$, and $F'_{i+1} \subseteq \sum_{j=\frac{a_{i+1}}{10(i+1)}}^{10(i+1)a_{i+1}} A(j) \otimes A(a_{i+1})$. Then there are homogeneous linear spaces $Z'_{i+1}, Z''_{i+1} \subseteq \sum_{j=\frac{a_{i+1}}{10(i+1)}}^{10(i+1)a_{i+1}} A(i)$ with $\dim_k Z'_{i+1} \leq \dim KF_{i+1}$, $\dim_k Z''_{i+1} \leq$*

$\dim KF_i'$ *and such that*

$$Q(F_{i+1}, a_{i+1}, a_{i+2}) + U \otimes A \subseteq U \otimes A + A \otimes M(Z'_{i+1}, a_{i+1}, a_{i+2}),$$

$$Q'(F'_{i+1}, a_{i+1}, a_{i+2}) + A \otimes U \subseteq A \otimes U + M(Z''_{i+1}, a_{i+1}, a_{i+2}) \otimes A$$

where $U = \sum_{k=1}^{\infty}(A(ka_{i+2} - a_{i+1}) + A(ka_{i+2}))U_{i+1}A$.

Proof. Using the fact that $U_{i+1} + Km_{i+1} = A(a_{i+1})$, it follows that for every $g \in F_{i+1}$ there is $z(g) \in \sum_{j=\frac{a_{i+1}}{10(i+1)}}^{10(i+1)a_{i+1}} A(j)$, and $\alpha \in K$ such that either $g \in \alpha m_{i+1} \otimes z(g) + U_{i+1} \otimes A$. Let $g_1, \ldots g_m$ be a basis of F_{i+1}. Let Z'_{i+1} be the linear space spanned by elements $z(g_1), z(g_2), \ldots, z(g_m)$. By the definition of $Q(F_{i+1}, a_{i+1}, a_{i+2})$ and of $M(Z_{i+1}, a_{i+1}, a_{i+2})$ it follows that $Q(F_{i+1}, a_{i+1}, a_{i+2}) + U \otimes A \subseteq U \otimes A + A \otimes M(Z'_{i+1}, a_{i+1}, a_{i+2})$. The proof that $Q'(F'_{i+1}, a_{i+1}, a_{i+2}) + A \otimes U \subseteq M(Z''_{i+1}, a_{i+1}, a_{i+2}) \otimes A + A \otimes U$ is similar. □

We are now ready to prove the main result of this chapter.

Theorem 6.1 *Suppose that we are given natural numbers* $0 < a_1 < a_2 < a_3 < \ldots$, *which satisfy assumptions of Lemma 5.3 and such that* $200(i+1)^2 a_i$ *divides* a_{i+1} *for every* i. *Suppose that for all* $i \geq 1$ *we are given homogeneous linear spaces* $G_i \subseteq A(a_i) \otimes A(a_i)$, $F_i \subseteq A(a_i) \otimes \sum_{j>\frac{a_i}{10i}}^{10ia_i} A(j)$, $F'_i \subseteq \sum_{j>\frac{a_i}{10i}}^{10a_i i} A(j) \otimes A(a_i)$ *with* $\dim_K G_i, \dim_K F_i, \dim_K F'_i < 2^{500ia_{i-1}} a_{i+2}^{100i}$, $F_1 = 0, G_1 = 0, G_2 = 0$. *Then there are linear spaces* U_i *and monomials* m_i *such that for all* $i \geq 1$ *we have*

1. $U_i \oplus Km_i = A(a_i)$ *and* $A(a_i) \cap \sum_{j=1}^{i-1} B_{a_{j+1}}(U_j) \subseteq U_i$,
2. $G(i) \subseteq U_i \otimes A(a_i) + A(a_i) \otimes U_i$,

Moreover there are homogeneous linear spaces $Z_i \subseteq \sum_{i=\frac{a_i}{10i}}^{10ia_i} A(i)$ *such that* $\dim_{Z_i} \leq \dim_K F_i + \dim_K F'_i$, *and such that*

3. $Q(F_j, a_j, a_{j+1}), Q'(F'_j, a_j, a_{j+1}) \subseteq D_i \otimes A + A \otimes D_i$ *where* $D_i = \sum_{j=1}^{i} B_{a_{j+1}}(U_j) + M(Z_j, a_j, a_{j+1})$.
4. $A(a_i) \cap D_{i-1} \subseteq U_i$.

Proof. We proceed by induction on i. For $i = 1$, $F_1, G_1 = 0$ so we can take U_1 to be any linear subspace of $A(a_1)$ of codimension 1. Then for an appriopriate monomial m_1 we have $U_1 + Km_1 = A(a_1)$. Then we can set $Z_1 = 0$.

Suppose that the result holds for all numbers not exceeding i, we will show that it holds for $i+1$. Assume that we already constructed sets $U_1, \ldots, U_i$, and sets $Z_1, \ldots, Z_i$. We will construct set U_{i+1} and then set Z_{i+1}.

Let V be a linear subspace of $A(a_{i+1})$ such that $V \oplus (D_i \cap A(a_{i+1})) = A(a_{i+1})$.

We will first show that

$$\dim_K(D_i \cap A(a_{i+1})) < \dim_K A(a_{i+1}) - 4\dim_K G_{i+1} - 2.$$

We will prove an equivalent statement that $\dim_K V > 4\dim_K G_{i+1} + 2$.

Let $\delta_{i+1} : A(a_{i+1}) \to A(a_{i+1})$ be as in Lemma 4.1. Then $\delta_{i+1}(U') = 0$, where $U' = (\sum_{j=1}^{i} B_{a_{j+1}}(U_j)) \cap A(a_{i+1})$. Denote $M' = (\sum_{j=1}^{i} M(Z_j, a_j, a_{j+1})) \cap A(a_{i+1})$, so $D_i \cap A(a_{i+1}) = U' + M'$ Then $\delta_{i+1}(V) + \delta_{i+1}(M') = \delta_{i+1}(A(a_{i+1})) = A(a_{i+1} \cdot t_{i+1})$. It follows that $\dim_K V \geq \dim_K \delta_{i+1}(V) \geq \dim_K A(a_{i+1} \cdot t_{i+1}) - \dim_K \delta_{i+1}(M')$, where t_{i+1} is as in Lemma 4.1.

Fix $n \leq i$, and let $f \in Z_n$, and denote $F = M(f, a_n, a_{n+1})$. Observe that $F \subseteq D_{n+k}$ for every $k \geq 0$, hence $A(a_{n+k+1}) \cap F \subseteq A(a_{n+k+1}) \cap D_{n+k}$, for all $k \geq 0$. If $n+k+1 \leq i$, then by the inductive assumption $A(a_{n+1+k}) \cap D_{n+k} \subseteq U_{n+k+1}$, and so $A(a_{n+1+k}) \cap F \subseteq U_{n+k+1}$. Therefore f satisfies assumptions of Lemma 4.2. Therefore any element f from sets Z_n satisfy assumptions of Lemma 4.2. Let $f_1, f_2, \ldots, f_{k'}$ be a basis of Z_n, We can apply Lemma 4.2 several times for $f = f_1$, $f = f_2, \ldots, f = f_{k'}$. Let $E(Z_n) = \bigcup_{j=1}^{k'} E(f_j)$ where $E(f_j)$ is as in Lemma 4.2 applied for $f = f_j$, then the cardinality of $E(Z_n)$ does not exceede $2a_n^4 \cdot 2^{2a_{n-1}} \cdot \dim_K Z_n < 2^{505a_{n-1}} a_{n+2}^{110i}$. Moreover, if $i + k + 1 \leq i + 1$ then

$$\delta_{n+k+1}(F \cap A[a_{n+k+1}]) \subseteq < E(Z_n) >$$

(where $< E(Z_n) >$ is the ideal generated in A by elements from $E(Z_n)$). In particular we have $\delta_{i+1}(F' \cap A[a_{i+1}]) \subseteq \sum_{n=1}^{i} < E(Z_n) >$, where we denote $F' = \sum_{j=1}^{k'} M(f_j, a_n, a_{n+1})$. Such sets $E(Z_n)$ can be constructed for all $n \leq i$. It follows that $\delta_{i+1}(D_i \cap A[a_{i+1}]) \subseteq < E(Z_n) >$ (because $\delta_{i+1}(U') = 0$ by Lemma 4.1). Let J be the ideal of A generated by all these sets $E(Z_n)$ for all $1 \leq n \leq i$.

By Lemma 4.2, $E(Z_n) \subseteq \sum_{i > \frac{a_n}{40n}}^{20na_n} A(i)$. It follows that I is generated by elements from sets $E(Z_n) \subseteq \sum_{i > \frac{a_n}{40n}}^{20na_n} A(i)$ with the cardinality of $E(Z_n)$ less than $2^{505a_{n-1}} a_{n+2}^{110i}$ for $n = 1, 2, \ldots, i$. By Theorem 5.3,

$$\dim_K(A(t_{i+1}a_{i+1})/A(t_{i+1}a_{i+1}) \cap I) > 4\dim_K G_{i+1} + 2.$$

Therefore, we obtain $dim_K V \geq \dim_K A(a_{i+1} \cdot t_{i+1}) - \dim_K \delta_{i+1}(M') = \dim_K A(t_{n+1}a_{i+1})/A(t_{n+1}a_{i+1}) \cap J$. It follows that $\dim_K V > 4\dim_K G_{i+1} + 2$, and so $\dim_K(D_i \cap A(a_{i+1})) < \dim_K A(a_{i+1}) - 4\dim_K G_{i+1} - 2$.

We can now apply Lemma 6.1 for $m = i+1$ and for sets $T = A(a_{i+1}) \cap D_i, Q = G_{i+1}$, and obtain set F as in Lemma 6.1. By Lemma 6.1 we get $T \subseteq F$ and $G_{i+1} = Q \subseteq F \otimes A(a_{i+1}) + A(a_{i+1}) \otimes F$, so we can set $U_{i+1} = F$.

We have constructed set U_{i+1} satisfying the thesis of our theorem. We can now put $Z_{i+1} = Z'_{i+1} + Z''_{i+1}$ where Z'_{i+1}, Z''_{i+1} are as in Lemma 6.2. □

Corollary 6.1 *Let $a_0, a_1, \ldots$ satisfy assumptions of Lemma 5.3 and Theorem 6.1. Let U_i, Z_i be as in Theorem 6.1, and let N be as in Theorem 3.1. Then $N \subseteq D \otimes A + A \otimes D$ where $D = \sum_{i=1}^{\infty} D_i$ with $D_i = \sum_{j=1}^{i} B_{a_{j+1}}(U_j) + M(Z_j, a_j, a_{j+1})$. Moreover, $A(a_i) \cap D_{i-1} \subseteq U_i$, for all $i \geq 1$.*

Moreover $S_i \subseteq A \otimes D_{i+3} + D_{i+3} \otimes A$ for every $i \geq 1$, where S_i is as in Theorem 3.1.

Proof. By Theorem 6.1, applied for sets F_i, F'_i, G_i as in Theorem 3.1, we have $A(a_i) \cap D_{i-1} \subseteq U_i$, for all $i \geq 1$. By Theorem 3.1, $N \subseteq \sum_{n=1}^{\infty} S_n$, where $S_n = \sum_{n=1}^{\infty} M_n + Q_n + Q_{n-1}$ and where $M_n = \sum_{0 \leq k,k'} ((A(ka_{n+1}) + (A(ka_{n+1} - a_n)) \otimes (A(k'a_{n+1}) + A(k'a_{n+1} - a_n)))G_n(A \otimes A)$,

$$Q_n = Q(F_n, a_n, a_{n+1}) + Q(F'_n, a_n, a_{n+1}).$$

By Lemma 6.2, $Q_n \subseteq D \otimes A + A \otimes D$. By Theorem 6.1, $G_i \subseteq U_i \otimes A + A \otimes U_i$, hence $M_n \subseteq D \otimes A + A \otimes D$. It follows that $N \subseteq A \otimes D + D \otimes A$.

Similarly, to show that $S_i \subseteq A \otimes D_{i+3} + D_{i+3} \otimes A$ observe that $S_i = M_n + Q_n + Q_{n-1}$. By Theorem 6.1 (2), and Lemma 6.2, we have $M_n \subseteq D_{i+3} \otimes A + A \otimes D_{i+3}$ because $\sum_{j=1}^{i} B_{a_{j+1}}(U_j) \subseteq D_i$. By Lemma 6.2 and by Theorem 6.1 (3), $Q_n, Q_{n-1} \in D_{n+3} \otimes A + A \otimes D_{n+3}$. □

Corollary 6.2 *Let notation be as in Corollary 6.1. Then there are homogeneous ideals $I_1, I_2, \ldots$, in $A \otimes A$ and such that $N = I_1 + I_2 + \ldots$, and for every n, $I_n \subseteq D_n \otimes A + A \otimes D_n$. Moreover, for every $f \in A \otimes A$ there is $n = n(f)$ such that $f^n, \xi(f^n) \in N'$ where $N' = \sum_{i=1}^{\infty} (A^{a_{n+2}} \otimes A^{a_{n+2}}) I_n$.*

Proof. Recall that N is constructed as in Theorem 3.1, hence for every $f \in A \otimes A$ there is $n = n(f)$ such that $f^n, \xi(f^n) \in N$. We put $I_1 = I_2 = \ldots = I_9 = 0$. If K is a countable field then for $n \geq 1$, we define I_{n+9} to be the ideal of $A \otimes A$ generated by elements $f_n^{20na_{n+1}}$ and $\xi(f_n^{20na_{n+1}})$ where f_n is as in the beginning of the proof of Theorem 3.1. If K is an uncountable field then for $n \geq 1$, we define for every $n \geq 1$, I_{n+9} be the ideal of $A \otimes A$ generated by elements from the set $\{f^{100a_{n+2}} : f \in \bar{H}_n\}$ and from the set $\{\xi(f)^{100a_{n+2}} : f \in \bar{H}_n\}$. From the proof of Theorem 3.1 it follows that $I_{n+9} \subseteq \sum_{i=1}^{n+5} S_i$, where S_i are as in Theorem 3.1. From the last part of Corollary 6.1, $S_n \subseteq D_{n+3} \otimes A + A \otimes D_{n+3}$ for every $n \geq 1$. Since $D_i \subseteq D_{i+1}$ for every n, it follows that $I_n \subseteq D_n \otimes A + A \otimes D_n$.

We will now show that for every $f \in A \otimes A$ there is $n = n(f)$ such that $f^n, \xi(f^n) \in N'$. Let $f \in N$, then by construction of ideals $I_1, I_2, \ldots$ there is m such that $f^n, \xi(f^n) \in I_m$ for some n. Then $f^{n+a_{m+2}} = f^{a_{m+2}} \cdot f^n \subseteq (A^{a_{m+2}} \otimes A^{a_{m+2}})I_m \subseteq N'$.

Observe also that $\xi(f^{n+2a_{m+2}}) = \xi(f^{a_{m+2}} \cdot f^n \cdot f^{a_{m+2}}) \subseteq (A^{a_{m+2}} \otimes A^{a_{m+2}})I_m \subseteq N'$. □

7 Definition of the ideal I

Let C be a right ideal of A, so $IA \subseteq A$, which is homogeneous. Define subspace $I[C]$ of A in the following way.

$$\{I[C] = \{r \in C : Ar \subseteq C\}.$$

Clearly $I(C)$ is an ideal in A.

Let $D = \bigcup_{i=1}^{\infty} D_i$, with $D_i = \sum_{j=1}^{i} B_{a_{j+1}}(U_j) + M(Z_j, a_j, a_{j+1})$, be as in Corollary 6.1. We can define $I[D] = \{a \in D : Aa \subseteq D\}$. As D is a right ideal, so $DA \subseteq D$, so it is clear that $I[D]$ is an ideal in D.

Lemma 7.1 *Let notations be as in Corollary 6.1 and Theorem 6.1. If* $a \in \bigcap_{i=0}^{a_{n+1}} A(i)D_n$ *then* $a \in I[D]$.

Proof. By the definition of D_n, $A(a_{n+1})D_n \subseteq D_n$. Let $v \in M$, we will show that $va \in D_n$, and hence $va \in D$. Let $v = v_1 v_2$ where the degree of v_1 divisible by a_{n+1}, $v_2 \in M(s)$ with $s < a_{n+1}$. Notice that by assumption $a \in A(a_{n+1} - s)D_n$ hence $v_2 a \in v_2 A(a_{n+1} - s)D_n \subseteq A(n+1)D_n \subseteq D_n$. Therefore $v_1 v_2 a \in v_1 D_n \subseteq D_n \subseteq D$.

So for every $v \in M$, $va \in D$, hence $Aa \subseteq D$, so $a \in I[D]$. □

Lemma 7.2 *Let notations be as in Corollary 6.1 and Theorem 6.1. Then* $A/I[D]$ *is an infinitely dimensional space.*

Proof. As $I(D) \subseteq D$, it suffices to show that A/D is an infinitely dimensional linear space. By the definition $D = \sum_{j=1}^{\infty} C_j$, where for every j, $C_j = B_{a_{j+1}}(U_j) + M(Z_j, a_j, a_{j+1})$ is a homogeneous linear space and all elements from C_j have degrees $\frac{a_j}{40j}$ or larger (elements with the smallest possible degrees comming from sets Z_j or U_j.) Suppose on the contrary that A/D is finitely dimensional, then it is nilpotent of degree j, for some j. Then, $A(j) \subseteq A(j) \cap \sum_{i=1}^{j} C_i$, so $A(j) \subseteq A \cap D_j$. As D_j is a right ideal it follows that $A(j+k) \subseteq D_j A(k) \subseteq D_j$ for all $k \geq 0$. Therefore, $A(a_{j+k}) \cap D_j = A(j+k)$ for all $k \geq 0$. Observe that for $k = 2$, $D_j \subseteq D_{j+1}$ and

by Corollary 6.1 we get $A(a_{j+2}) \cap D_j \subseteq A(a_{j+2}) \cap D_{j+1} \subseteq U_{j+2} \neq A(a_{j+2})$, a contradiction. □

We will now prove two supporting Lemmas.

Lemma 7.3 *Let $Q_1, Q_2, B \subseteq A$, are linear spaces. If $a \in A \otimes Q_1 + B \otimes A$ and $a \in A \otimes Q_2 + B \otimes A$ then $a \in A \otimes (Q_1 \cap Q_2) + B \otimes A$. Similarly, if $a' \in Q_1 \otimes A + A \otimes B$ and $a' \in Q_2 \otimes A + A \otimes B$, then $a' \in (Q_1 \cap Q_2) \otimes A + A \otimes B$.*

Proof. If we take $B' = A/B$ then $a + B \in B' \otimes Q_1, a + B \in B' \otimes Q_2$, hence $a + B \in B' \otimes (Q_1 \cap Q_2)$, hence $a \in A \otimes (Q_1 \cap Q_2) + B \otimes A$. The proof of the second statement is similar. □

Lemma 7.4 *Let n be a natural number, and let $Q_1, \ldots, Q_n$ be linear subspaces of A, and let $a \in A \otimes A$ be such that $a \in Q_i \otimes A + A \otimes Q_j$ for all $i, j \leq n$. Then $a \in (\bigcap_{i=1}^n Q_i) \otimes A + A \otimes (\bigcap_{i=1}^n Q_i)$.*

Proof. Observe that by the first part of Lemma 7.3 applied several times (for sets $Q_1, \ldots Q_n$) we get, that for every $j \leq n$, $a \in A \otimes \bigcap_{i=1}^n Q_i + Q_j \otimes A$. By the second part of Lemma 7.3 applied to $a' = a$ we get $a \in (\bigcap_{i=1}^n Q_i) \otimes A + A \otimes (\bigcap_{i=1}^n Q_i)$. □

Lemma 7.5 *Let notation be as in Corollary 6.1 and Theorem 6.1. Let I_n be a homogeneous ideal of $A \otimes A$ such that $I_n \subseteq A \otimes D_n + D_n \otimes A$ for some n. Then $(A^{a_{n+2}} \otimes A^{a_{n+2}}) I_n \subseteq A \otimes I[D] + I[D] \otimes A$.*

Proof. We will first show that $(A^{a_{n+2}} \otimes A^{a_{n+2}}) I_n$ is contained in $A(k) D_n \otimes A + A \otimes A(j) D_n$ for each $0 \leq i, j \leq a_{n+1}$. Note that $(A^{a_{n+2}} \otimes A^{a_{n+2}}) I_n = (A(k) \otimes A(j))(A^{a_{n+2}-k} \otimes A^{a_{n+2}-j}) I_n \subseteq (A(k) \otimes A(j)) I_n \subseteq (A(k) \otimes A(j))(D_n \otimes A + A \otimes D_n) \subseteq A(k) D_n \otimes A + A \otimes A(j) D_n$.

Let $a \in A^{a_{n+2}} I_n$. Then a satisfies assumptions of Lemma 7.4 for $Q_i = A(i) D_n$. By Lemma 7.4, $a \in A \otimes P = P \otimes A$ where $P = \bigcap_{i=0}^{a_{n+1}} A(i) D_n$. By Lemma 7.1, $P \subseteq I[D]$. Hence $a \in D_n \otimes A + A \otimes A(j) D_n$. □

8 Proof of Theorem 11.2

We are now ready to prove Theorem 11.2.

Proof of Theorem 11.2 Let notation be as in Corollary 6.1, Corollary 6.2 and Theorem 3.1. Let $D = \sum_{i=1}^{\infty} D_i$ as in Corollary 6.1 and let $I[D] = \{r \in D : Ar \subseteq D\}$. Let $R = A/I[D]$, we claim that $R \otimes R$ and $R \otimes R^{op}$ are nil algebras.

By Corollary 6.2 there are homogeneous ideals $I_1, I_2, \ldots,$ in $A \otimes A$ and such that $N = I_1 + I_2 + \ldots$, and for every n, $I_n \subseteq D_n \otimes A + A \otimes D_n$.

Moreover, for every $f \in A \otimes A$ there is $n = n(f)$ such that $f^n, \xi(f^n) \in N'$, where $N' = \sum_{i=1}^{\infty}(A^{a_{n+2}} \otimes A^{a_{n+2}})I_n$. Observe that by Lemma 7.5, $(A^{a_{n+2}} \otimes A^{a_{n+2}})I_n \subseteq A \otimes I[D] + I[D] \otimes A$. Therefore $N' \subseteq A \otimes I[D] + I[D] \otimes A$. By Assumption for every $f \in A \otimes A$ there is number $n = n(f)$ such that $f^n \in N' \subseteq A \otimes I[D] + I[D] \otimes A$. It follows that $(A/I) \otimes_K (A/I) = R \otimes_K R$ is a nil algebra.

Let $g \in A \otimes A$, and let $f \in A \otimes A$ be such that $\xi(f) = g$ in the linear space $A \otimes A$. By assumption there is number $n = n(f)$ such that $\xi(f^n) \in N' \subseteq A \otimes I[D] + I[D] \otimes A$.

Let $*$ be the opposite multiplication. Let $g' = g * g * \ldots * g$ -the opposite multiplication of m copies of g. Then in the linear space $A \otimes A$ we have $g' = \xi(f^m)$. By Assumption $\xi(f^n) \in N'$ hence $g' \subseteq J$ where $J = A \otimes I[D] + I[D] \otimes A$. J is a linear subspace in $A \otimes A$, so $g' + J = 0$ in $A/I[D] \otimes A/I[D]$. It follows that $(g + J) * (g + J) * \ldots * (g + J) \subseteq g' + J$, so it follows that $g + J$ is a nilpotent element in $A/I[D] \otimes (A/I[D])^{op}$ for all $g \in A$. Therefore, $R \otimes R^{op}$ is a nil algebra.

9 Appendix: Proof of Theorem 3.1. Nil ideals

In this section we will recall some ideas from [21]. Results from [21] were simplified in [14], so we will use [14]. We first introduce the following notation: let $r \in (A(1) + \ldots + A(t)) \otimes (A(1)) + \ldots + A(q))$, where t, q are the minimal possible. Then we say that the left degree of a, denoted $l(a)$ is t, and the right degree of a denoted $r(a)$ is q.

We begin with three lemmas which correspond to Lemma 1 from [21] and Lemma 7 from [14].

Lemma 9.1 *Let $f \in \sum_{1 \leq i,j \leq t} A(i) \otimes A(j)$ for some $t \geq 1$. For each $n \geq t$ there exists a linear space $F(f, n) \subseteq A(n) \otimes A$ with*

$$\dim_K F(f, n) < 2^{2t+1} tn$$

such that, for every integer $m \geq n$, f^m belongs to the right ideal of $A \otimes A$ generated by polynomials of the form hgf^{m-n}, where $h \in F(f, n)$ and $g \in (A(0) + \ldots + A(t)) \otimes K$.

Proof. We will modify, in a straightforward way, the proof of Lemma 7 in [14], when we only consider the first component in $A \otimes A$. We write $f = \sum_{i=1}^{t} f_i$, where $f_i \in A(i) \otimes A$; thus f^m is a sum of terms of the

form $f_{i_1}f_{i_2}\dots f_{i_k}f_{i_{k+1}}\dots f_{i_m}$, where k denotes the largest subscript such that $i_1+i_2+\ +i_k\leq n$. We then have the decomposition:

$$f^m=\sum_{k\leq n}\sum_{i_1+\dots+i_k=n} f_{i_1}\dots f_{i_k}f^{m-k}+\sum_{j=1}^{t-1}(\sum_{i_1+\dots+i_k=n-j,i_{k+1}>j} f_{i_1}\dots f_{i_k}f_{i_{k+1}}f^{m-k-1}))$$

Hence

$$f^n=\sum_{k\leq n}v_{0,k}f^{m-k}+\sum_{k\leq n-1}(\sum_{j=1}^{t-1}v_{j,k}(f_{j+1}+f_{j+2}+\dots+f_t)f^{m-k-1})$$

where $v_{j,k}$ is the homogeneous component of degree $n-j$ in f^k with respect to the first component in the tensor product (that means that $f^k=\sum_j v_{j,k}$ and $v_{j,k}\in A(n-j)\otimes A$, for each j).

Taking $F(f,n)=\sum_{i=0}^{t}\sum_{j=0}^{t-1}\sum_{k=1}^{n}v_{j,k}(M(j)\otimes A(i))$, the representation above has the desired properties (where K is the base field). To see this, note that in the second summation each term $v_{j,k}f_{j+i}f^{m-k-1}$ lies in the span of all members of $v_{j,k}(M(j+i)\otimes\sum_{j'=1}^{t}M(j'))(M(i))f^{m-n}f^{n-k-1}\subseteq v_{j,k}(M(j)\otimes\sum_{j'=1}^{t}M(j'))(M(i)\otimes K)f^{m-n}(A\otimes A)$. As elements of $v_{j,k}(M(j)\otimes\sum_{j'=1}^{t}M(j'))$ lie in $F(f,n)$, and elements of $M(i)$ have degree $<t$, the product resides in the desired right ideal. The same is still more easily seen for terms of the first summation. Notice that if we want $v_{j,k}$ to be nonzero, then we necessarily must have $\frac{n-j}{t}\leq k\leq n-j$. We compute

$$\dim_K G(f,n)<\sum_{j=0}^{t-1}\sum_{k\geq\frac{(n-j)}{t}}^{n-j}2^t2^{t+1}\leq 2^{2t+1}tn$$

□

Let i,j be nonnegative integers. For $a\in A(i)\otimes A(j)$ we denote the right degree of a as $r(a)$, and the left degree of a by $l(a)$. Here, $l(a)=i$, $r(a)=j$.

We will now give a generalization of this lemma.

Lemma 9.2 *Let $f\in\sum_{1\leq i,j\leq t}A(i)\otimes A(j)$ for some $t\geq 1$. For each $n>t$ there exists a linear space $F(f,n)\subseteq A(n)\otimes\sum_{i\geq\frac{n}{2t}}^{2tn}A(i)$ with $\dim_K F(f,n)\leq 2^{2t+1}tn$ such that, for every integer $m\geq n$, f^m belongs to the right ideal of $A\otimes A$ generated by elements of the form hgf^{m-n}, where $h\in F(f,n)$ and $g\in\sum_{0\leq i\leq t}A(i)\otimes K$.*

Proof. With the notation from the previous lemma, we have $f=\sum_{i=1}^{t}f_i$, where $f_i\in A(i)\otimes(A(1)+\dots+A(t))$. We can now write $f_i=\sum_{j=1}^{t}f_{i,j}$, where

$f_{i,j} \in A(i) \otimes A(j)$. Observe that for every i, j, $l(f_{i,j}) \leq t \cdot r(f_{i,j})$ and $r(f_{i,j}) \leq t \cdot l(f_{i,j})$, therefore $\frac{1}{t} \cdot l(f_{i,j}) \leq r(f_{i,j}) \leq t \cdot l(f_{i,j})$. Similarly if $0 \leq i, j \leq t$, then for an element $g \in f_{i,j}(A(i) \otimes A(j))$ we have $\frac{1}{2t} \cdot l(g) \leq r(g) \leq 2t \cdot l(g)$. It follows that if $m \in A \otimes A$ is a product of some elements $f_{i,j}$ or such elements g then $\frac{1}{2t} \cdot l(m) \leq r(m) \leq 2t \cdot l(m)$. By the definition of the set $F(f, n)$ from the previous Lemma, and because $n > t$, we get that any element from $F(f, n)$ there is a sum of elements g with $\frac{l(g)}{2t} \leq r(g) \leq 2t \cdot l(g)$. By the previous lemma $F(f, n) \subseteq A(n) \otimes A$ and hence $F(f, n) \subseteq A(n) \otimes \sum_{i \geq \frac{n}{2t}}^{2tn} A(i)$, as required. Also by construction of element g in thesis of Lemma 9.1 we have $g \in \sum_{0 \leq i \leq t} A(i) \otimes K$, as required. □

Lemma 9.3 *Let $f \in H$ where $H = \sum_{1 \leq i,j \leq t} A(i) \otimes A(j)$ for some $t \geq 0$, and let $n > 2t$. Let $F(f, r)$ be as in Lemma 9.2. Let $T(f, n)$ be the smallest homogeneous linear space containing $\bigcup_{0 \leq i,j \leq t}(A(i) \otimes A(j))F(f, n - i)$. Then $T(f, n) \subseteq A(n) \otimes \sum_{i \geq \frac{n-t}{2t}}^{2tn+t} A(i)$ and $\dim_K T(f, n) < 2^{4t+3} n^3 t^2$. Moreover, for every $m > n$, $f^m, H' f^m \subseteq T(f, n) H' f^{m-n}(A \otimes A)$, where $H' = \sum_{0 \leq i,j \leq t} A(i) \otimes A(j)$.*

Proof. By Lemma 9.2, $F(f, n - i) \subseteq A(n - i) \otimes \sum_{i \geq \frac{n-i}{2t}}^{2t(n-i)} A(i)$. Hence, $F(f, n-i)(A(i) \otimes A(j)) \subseteq A(n) \otimes \sum_{i \geq \frac{n-t}{2t}}^{2tn+t} A(i)$. By the definition of $T(f, n)$ we get $T(f, n) \subseteq A(n) \otimes \sum_{i \geq \frac{n-t}{2t}}^{2tn+t} A(i)$.

By Lemma 9.2, $\dim_K F(f, n - i) < 2^{2t+1} tn$, also $\dim_K A(i) = 2^i$, hence $\dim_K T(f, n) < 2^{2t+1} tn \cdot 2^{2t+2} \cdot (2tn + t) \leq 2^{4t+3} t^2 n^3$ (since $n > 2t \geq 2$).

Let $i, i' \leq t$. Observe now that by Lemma 9.2 applied for $n - i$, $(A(i) \otimes A(i'))f^m \subseteq (A(i) \otimes A(i'))F(n-i)H' f^{m-n+i}(A \otimes A) \subseteq T(f, n)H' f^{m-n+i}(A \otimes A)$, as required. □

The following lemmas can be proven in the same way as Lemma 9.2, when we interchange the left side with the right side.

Lemma 9.4 *Let $f \in \sum_{1 \leq i,j \leq t} A(i) \otimes A(j)$ for some $t \geq 1$. For each $n > t$ there exists a linear space $F'(f, n) \subseteq \sum_{i \geq \frac{n}{2t}}^{2tn} A(i) \otimes A(n)$ with $\dim_K F'(f, n) \leq 2^{2t+1} tn$ such that, for every integer $m \geq n$, f^m belongs to the right ideal of $A \otimes A$ generated by elements of the form hgf^{m-n}, where $h \in F'(f, n)$ and $g \in K \otimes \sum_{0 \leq i \leq t} A(i)$.*

Lemma 9.5 *Let $f \in H$ where $H = \sum_{1 \leq i,j \leq t} A(i) \otimes A(j)$ for some $t \geq 0$, and let $n > 2t$. Let $F'(f, r)$ be as in Lemma 9.4. Let $T'(f, n)$ be the smallest homogeneous linear space containing $\bigcup_{0 \leq i,j \leq t}(A(i) \otimes A(j))F'(f, n-j)$. Then $T'(f, n) \subseteq \sum_{i \geq \frac{n-t}{2t}}^{2tn+t} A(i) \otimes A(n)$ and $\dim_K T'(f, n) < 2^{4t+3} n^3 t^2$. Moreover,*

for all n, we have for every integer $m \geq n$, $H'f^m \subseteq T'(f,n)H'f^{m-n}$, where $H' = \sum_{0\leq i,i'\leq t} A(i)\otimes A(i')$.

Lemma 9.6 *Let $f \in H$ where $H = \sum_{1\leq i,j\leq t} A(i)\otimes A(j)$ for some $t \geq 0$, and let $n \leq 2t$. Define $T'(f,n) = \sum_{i\geq n-t}^{2tn} A(i)\otimes A(n)$. Then $\dim_K T'(f,n) < 2^{2tn+1} \leq 2^{2t^2+1}$. Moreover, for all $n \leq 2t$, and for every integer $m \geq n$, $H'f^m \in T'(f,n)H'f^{m-n}(A\otimes A)$, where $H' = \sum_{0\leq i,j\leq t} A(i)\otimes A(j)$.*

Proof. We proceed by induction on n, let $n = 1$. If $1 \leq i'$, then $(A(i)\otimes A(i'))f^m \subseteq (A(i)\otimes A(1))(K\otimes A(i'-1))f^m \subseteq T(f,1)H'f^m$. If $i' = 0$ then $(A(i)\otimes K)f^m = ((A(i)\otimes K)f)\cdot f^{m-1} \subseteq (\sum_{i=0}^{2t} A(i)\otimes A(1))(K\otimes \sum_{i<t} A(i))f^{m-1} \subseteq T(f,1)H'f^m$. We now assume the result is true for some number n (and all numbers $m \leq n$), and prove it for $n+1$.

Let $i' > 0$ then $(A(i)\otimes A(i'))f^m \subseteq (A(i)\otimes A(1))(K\otimes A(i'-1))f^m$. By the inductive assumption, $(A(i)\otimes A(i'-1))f^m \subseteq T'(f,n)H'f^{m-n}(A\otimes A)$. Hence $(A(i)\otimes A(i'))f^m \subseteq (A(i)\otimes A(1))T'(f,n)H'f^{m-n}(A\otimes A) \subseteq T'(f,n+1)H'f^{m-n}(A\otimes A)$.

If $i' = 0$, then similarly as before $(A(i)\otimes A(i'))f^m \subseteq ((A(i)\otimes K)f)\cdot f^{m-1} \subseteq (\sum_{j=1}^{2t} A(j)\otimes A(1))H'f^{m-1}$. By the inductive assumption $H'f^{m-1} \subseteq T'(f,n)$. Notice that $(\sum_{j=1}^{2t} A(j)\otimes A(1))T(f,n) \subseteq T'(f,n+1)$, so $(A(i)\otimes A(i'))f^m \subseteq T'(f,n+1)H'f^{m-1}$, as required □

10 Appendix: Proof of Theorem 3.1. Linear spaces G_1 and G_2

We now introduce a simple lemma which can be read independently from the rest of this paper.

Lemma 10.1 *Let α, t, c be a natural numbers with $c \leq \alpha$, and let G_1, G_2 be homogeneous linear subspaces with*

$$G_1 \subseteq \sum_{i=\alpha}^{\alpha+t} A(i)\otimes \sum_{i=c}^{\alpha+t} A(i), G_2 \subseteq \sum_{i=c}^{\alpha+t} A(i)\otimes \sum_{i=\alpha}^{\alpha+t} A(i).$$

Then there exist linear spaces

$$T_\alpha(G_1) \subseteq A(\alpha)\otimes \sum_{i=c}^{\alpha} A(i), T_\alpha(G_2) \subseteq \sum_{i=c}^{\alpha} A(i)\otimes A(\alpha),$$

and such that

$$G_1 \subseteq T_\alpha(G_1)(\sum_{0\leq i,j\leq t} A(i)\otimes A(j)), G_2 \subseteq T_\alpha(G_2)(\sum_{0\leq i,j\leq t} A(i)\otimes A(j)).$$

Moreover, $\dim_K T_\alpha(G_1) \leq 2^{2t+2}\dim_K G_1$ *and* $\dim_K T_\alpha(G_2) \leq 2^{2t+2}\dim_K G_2$.

Proof. We will do a proof for the set G_1; proof for the set G_2 is similar. Let $c_1, c_2, \ldots, c_\gamma$ be a basis of $\sum_{0\leq i,j\leq t} A(i)\otimes A(j)$. Notice that $\gamma \leq 2^{2t+2}$, because $\dim_k A(i) = 2^i$. Then we can write every element $g \in G_1$ in the form $g = \sum_{i=c}^{\gamma} g_i c_i$, where $g_i \in A(\alpha)\otimes \sum_{c\leq i\leq \alpha} A(i)$. Let $g(1), g(2), \ldots$ be a basis of the linear space G. We can do such a decomposition $g(k) = \sum_{i=1}^{\gamma} g(k)_i c_i$ for every element from the basis of G. We can now put $T_\alpha(G_1)$ to be a linear space spanned by all these elements $g(k)_i$; it is clear that set $T_\alpha(G_1)$ satisfies the thesis of the theorem. □

We will now prove another modification of Lemma 1 from [21], and Lemma 7 from [14].

Lemma 10.2 *Let* $f \in \sum_{1\leq i,j\leq t} A(i)\otimes A(j)$ *for some* $t \geq 1$. *For each* $n > t$ *there exist homogeneous linear spaces* $G_1(f,n) \subseteq A(n)\otimes \sum_{i\geq \frac{n}{2t}}^{n} A(i)$, $G_2(f,n) \subseteq \sum_{i\geq \frac{n}{2t}}^{n} A(i)\otimes A(n)$ *with* $\dim_K G_1(f,n), \dim_K G_2(f,n) \leq 2^{2t+6}n^2t^3$ *such that, for every integer* $m \geq n+1$, f^m *belongs to the right ideal of* $A\otimes A$ *generated by polynomials of the form* hgf^{m-n-1}, *where* $h \in G_1(f,n)\bigcup G_2(f,n)$ *and* $g \in \sum_{0\leq i,j\leq t} A(i)\otimes A(j)$.

Proof. Write $f = \sum_{1\leq i,j\leq t} f_{i,j}$ where $f_i \in A(i)\otimes A(j)$. Then f^m is a sum of terms of the form

$$f_{i_1,j_1}f_{i_2,j_2}\ldots f_{i_k,j_k}f_{i_{k+1},j_{k+1}}\ldots f_{i_m,j_m}$$

where k denotes the largest subscript such that both $i_1+i_2+\ldots+i_k \leq n$ and $j_1+j_2+\ldots+j_k \leq n$. We then have the decomposition:

$$f^m = \sum_{k\leq n}[(q_k+q'_k+q''_k)f^{m-k} + (\sum_{0\leq i,j<t} q_{k,i,j} + \sum_{0\leq i<t} l_{k,i} + \sum_{0\leq i<t} \bar{l}_{k,i})f^{m-k-1}]$$

where

$$q_k = \sum_{i_1+\ldots+i_k=n; j_1+\ldots+j_k=n} f_{i_1,j_1}\ldots f_{i_k,j_k},$$

$$q'_k = \sum_{i_1+\ldots+i_k=n; j_1+\ldots+j_k<n} f_{i_1,j_1}\ldots f_{i_k,j_k},$$

$$q''_k = \sum_{i_1+\ldots+i_k<n; j_1+\ldots+j_k=n} f_{i_1,j_1}\ldots f_{i_k,j_k},$$

$$q_{k,i,j} = \sum_{i_1+...+i_k=n-i, i_{k+1}>i; j_1+...+j_k=n-j, j_{k+1}>j} f_{i_1,j_1} \dots f_{i_k,j_k} f_{i_{k+1},j_{k+1}}$$

$$l_{k,i,} = \sum_{i_1+...+i_k=n-i, i_{k+1}>i; j_1+...+j_k+j_{k+1}\leq n} f_{i_1,j_1} \dots f_{i_k,j_k} f_{i_{k+1},j_{k+1}}$$

$$\bar{l}_{k,j} = \sum_{j_1+...+j_k=n-i, j_{k+1}>i; i_1+...+i_k+i_{k+1}\leq n} f_{i_1,j_1} \dots f_{i_k,j_k} f_{i_{k+1},j_{k+1}}$$

Notice that, similarly to Lemma 9.2, we get that

$$q, q', q'', q_{k,i,j}, l_{k,i}, \bar{l}_{k,i} \subseteq (\sum_{i=0}^{t} A(n+i) \otimes \sum_{i\geq \frac{n}{2t}}^{2tn} A(i)) + (\sum_{i\geq \frac{n}{2t}}^{2tn} A(i) \otimes \sum_{i=0}^{t} A(n+i)).$$

On the other hand, by construction $q, q', q'', q_{k,i,j}, l_{k,i}, \bar{l}_{k,i} \subseteq \sum_{i,j\leq n+t} A(i) \otimes A(j)$. It follows that

$$q, q', q'', q_{k,i,j}, l_{k,i}, \bar{l}_{k,i} \subseteq (\sum_{i=0}^{t} A(n+i) \otimes \sum_{i\geq \frac{n}{2t}}^{n+t} A(i)) + (\sum_{i\geq \frac{n}{2t}}^{n+t} A(i) \otimes \sum_{i=0}^{t} A(n+i)).$$

Let G be a homogeneous linear space spanned by all homogeneous components of q_k, q'_k, q''_k, $q_{k,i,j}$, $l_{k,i}$, $\bar{l}_{k,i}$, then $dim_K G \leq 6nt^2 \cdot (n+t)t = 6n(n+t)t^3$. Let c be the smallest natural number which equals at least $\frac{n}{2t}$ and denote $\alpha = n$. Then there are linear spaces

$$G_1 \subseteq (\sum_{i=\alpha}^{\alpha+t} A_i \otimes \sum_{i=c}^{\alpha+t} A_i), G_2 \subseteq (\sum_{i=c}^{\alpha+t} A_i \otimes \sum_{i=\alpha}^{\alpha+t} A_i)$$

such that $G = G_1 + G_2$ and $\dim_K G_1, \dim_K G_2 \leq \dim_K G$. We can now take $G_1(f,n) = T_n(G_1)$, $G_2(f,n) = T_n(G_2)$ as in the Lemma 10.1 applied for $\alpha = n$. By Lemma 10.1, $\dim_K G_1(f,n), \dim_K G_2(f,n) \leq 2^{2t+2} \dim_K G \leq 2^{2t+2} 6n(n+t)t^3 < 2^{2t+6} n^2 t^3$, as required. □

11 Appendix: Proof of Theorem 3.1. Introducing sets $Q(S,n,m)$

Before introducing sets $Q(S,n,m)$, we will prove the following lemma.

Lemma 11.1 *Let $f \in H$ where $H = \sum_{1\le i,j\le t} A(i)\otimes A(j)$ for some $t \ge 1$. Let $n_1 > t$, $n_2 - 3n_1 > t$, $n_3 > 2tn_2$, $m > 8n_3 + 6tn_1$ be natural numbers. Let $G_1(f,n)$, $G_2(f,n)$ be as in Lemma 10.2, and $T(f,n)$, $T'(f,n)$ be as in Lemmas 9.3 , 9.5 and 9.6. Then the two-sided ideal of $A\otimes A$ generated by f^m belongs to the right ideal of $A\otimes A$ generated by elements from*

$$\sum_{1\le k,a_2<j} ((A(k\cdot n_3-n_2)\otimes A(j))T(f,n_2-n_1)T(f,n_1)(E(f,n_1,n_2)\cap E'(f,n_1,n_2))$$

where $E(f,n_1,n_2) = T(f,n_1)T(f,n_2-n_1)(A\otimes A)$, $E'(f,n_1,n_2) = S(f,n_1,n_2)+ S'(f,n_1,n_2)$,

$$S(f,n_1,n_2) = G_1(f,n_2-n_1)T(f,n_1)(A\otimes A)$$

$$S'(f,n_1,n_2) = G_2(f,n_2-n_1)\bigcap_{i=0}^{2n_1}(T'(f,i)T'(f,n_1)(A\otimes A)).$$

Moreover, linear spaces $E(f,n_1,n_2)$, $E'(f,n_1,n_2)$, $S(f,n_1,n_2)$, $S'(f,n_1,n_2)$ are homogeneous.

Proof. The ideal generated by f^m is spanned by elements $(A(k\cdot n_3 - n_2 - v)\otimes A(j))f^m$, for some $n_3 \le v < 2n_3$, and some $0 \le k,j$. By Lemma 9.2, f^m belongs to the right ideal of $A\otimes A$ generated by elements of the form hgf^{m-v}, where $h \in F(f,v)$ and $g \in \sum_{0\le i\le t} A(i)\otimes K$. Recall that $\frac{v}{t} \ge \frac{n_3}{t} > a_2$. Therefore, the ideal generated by f^m belongs to the right ideal generated by elements from $\sum_{1\le k,n_2\le i}((A(k\cdot n_3 - n_2)\otimes A(i))Hf^{m-v}(A\otimes A)$ where $H = \sum_{0\le i,j\le t} A(i)\otimes A(j)$. By Lemma 9.3, we get $Hf^{m-v} \subseteq T(f,n_2-n_1)Hf^{m-v-n_2+n_1}(A\otimes A)$. Applying Lemma 9.3 again, we get $Hf^{m-v-n_2+n_1} \subseteq T(f,n_1)Hf^{m-v-n_2}(A\otimes A)$. Therefore, the two sided ideal of $A\otimes A$ generated by f^m is contained in $\sum_{1\le,a_2<i}(A(k\cdot n_3-n_2)\otimes A(i))T(f,n_2-n_1)T(f,n_1)Hf^{m-n_2-v}(A\otimes A)$. Recall that $v \le 2n_3$, so it suffices to prove that $Hf^{m-n_2-2n_3} \subseteq E(f,n_1,n_2)\cap E'(f,n_1,n_2)$. By Lemma 9.3, applied for $n = n_1$ and then for $n = n_2 - n_1$ we get $Hf^{m-n_2-2n_3} \subseteq T(f,n_1)T(f,n_2-n_1)(A\otimes A)$, so $Hf^{m-n_2-2n_3} \subseteq E(f,n_1,n_2)$.

On the other hand, by Lemma 10.2, applied to $n = n_2 - n_1$ we get that $Hf^{m-n_2-2n_3} \subseteq (G_1(f,n_2-n_1)+G_2(f,n_2-n_1))Hf^{m-2n_2-2n_3+n_1-1}(A\otimes A)$.

Clearly, $m-2n_2-2n_3+n_1-1 \ge m-4n_3$. By Lemma 9.3, applied to $n = n_2$, we get $G_1(f,n_2-n_1)Hf^{m-4n_3} \subseteq G_1(f,n_2-n_1)T(f,n_1)(A\otimes A) = S(f,n_1,n_2)(A\otimes A) \subseteq E(f,n_1,n_2)$.

If $i > 2t$ then we apply Lemma 9.5 for $n = i$; if $i \le 2t$ then we apply Lemma 9.6 and we get $Hf^{m-4n_3} \subseteq T'(f,i)H'f^{m-4n_3-i}(A\otimes A)$ where $H' = \sum_{0\le l,l'\le t} A(l)\otimes A(l')$. We then apply 9.5 for $n = n_1$ to get $Hf^{m-4n_3} \subseteq$

$T'(f,i)T'(f,n_1)(A\otimes A)$ for every $n_1 \le i \le 2n_1$. Therefore $G_2(f,n_2-n_1)Hf^{m-4n_3} \subseteq G_2(f,n_2-n_1)\bigcap_{i=n_1}^{2n_1}(T'(f,i)T'(f,n_1)(A\otimes A)) \subseteq E'(f,n_1,n_2)$.

Moreover, sets $E(f,n_1,n_2)$, $E('f,n_1,n_2)$, $S(f,n_1,n_2$, $S'(f,n_1,n_2$ are homogeneous, because by Lemmas 9.3, 9.5 sets $T(f,n_1)$ and $T(f,n_2)$ are homogeneous. □

Let $1 \le \alpha$, $0 < n_1 < n_2$ be natural numbers and let $F \subseteq A(n_1)\otimes A(\alpha)$. Recall that

$$Q(F,n_1,n_2) = \sum_{k=0}^{\infty} \sum_{j\in W(n_1,n_2)} ((A(k\cdot n_2 - n_1) + A(k\cdot n_2))\otimes A(j))F(A\otimes A),$$

where $j \in W(n_1,n_2)$ if and only if the interval $[j,j+\alpha]$ is disjoint with all intervals $[kn_2-n_1, kn_2+n_1]$ for $k = 0,1,2,\ldots$.

The following lemma will be used in the proof of Lemma 11.3.

Lemma 11.2 *Given a natural number n, and a linear space $D \subseteq A \otimes \sum_{i=1}^{\infty} A(n+i)$ let $Z_n(D)$ to be the smallest linear space that $D \subseteq Z_n(D)(K\otimes A(n))$. Similarly, let $\bar{Z}_n(D)$ to be the smallest linear space that $D \subseteq (K\otimes A(n))\bar{Z}_n(D)$. Then $\dim_K Z_n(D), \dim_K \bar{Z}_n(D) \le 2^n \dim_K D$.*

Proof. Similar to Lemma 10.1. □

Lemma 11.3 *Let $f \in H$ where $H = \sum_{1\le i,j\le t} A(i)\otimes A(j)$ for some $t \ge 1$. Let $0 < n_1 < n_2 < n_3$ be natural numbers such that n_1 divides n_2, n_2 divides n_3 and $10tn_1 < n_2$, $10tn_2 < n_3$, $10tn_3 < m$, $10t < n_1$. Let $U(n_1,n_3) = \bigcup_{k=1}^{\infty}[kn_3-n_1, kn_3+n_1]$. Let $G_1(f,n)$, $G_2(f,n)$ be as in Lemma 10.2, and $T(f,n)$, $T'(f,n)$ be as in Lemmas 9.3, 9.5 and 9.6, and $E'(f,n_1,n_2)$ be as in Lemma 11.1. Then there is set $F(f,n_1,n_2,n_3) \subseteq A(n_2)\otimes \sum_{i=1}^{2tn_2+2t} A(i)$ with $\dim_K F(f,n_1,n_2,n_3) \le 2^{40tn_1}n_2$ and such that the two-sided ideal of $A\otimes A$ generated by f^m belongs to the right ideal of $A\otimes A$ generated by elements from $Q + P(A(n_3)\otimes A(n_3))$ where $Q = Q(T(f,n_1),n_1,n_2) + Q(F(f,n_1,n_2,n_3),n_2,n_3)$, and where*

$$P = \sum_{k=1}^{\infty} \sum_{j\in U(n_1,n_3)} (A(k\cdot n_3)\otimes A(j))E'(f,n_1,n_2)(A\otimes A).$$

Moreover, if $n_2 > 100tn_1$ then $F(f,n_1,n_2,n_3) \subseteq A(n_2)\otimes \sum_{i>\frac{n_2}{10t}}^{2tn_2+2t} A(i)$.

Proof. Let $D = T(f,n_2-n_1)T(f,n_1) + T(f,n_1)T(f,n_2-n_1) + T(f,n_2)$,

$$F(f,n_1,n_2,n_3) = D + Z_{5tn_1}(D) + \bar{Z}_{5tn_1}(D),$$

where $Z_n(D)$, $\bar{Z}_n(D)$ are as in Lemma 11.2. By Lemma 11.2, we get $\dim_K F(f,n_1,n_2,n_3) \leq (1+2^{5tn_1+1})\dim_K D \leq 2^{40tn_1}n_2$. By Lemma 9.3, $T(f,n) \subseteq A(n) \otimes \sum_{i>\frac{n-t}{2t}}^{2nt+t} A(i)$, so $F(f,n_1,n_2,n_3) \subseteq A(n_2) \otimes \sum_{i>\frac{n_2-2t}{2t}-5tn_1}^{2tn_2+2t} A(i)$. Notice that if $n_2 > 100tn_1$ then $F(f,n_1,n_2,n_3) \subseteq A(n_2) \otimes \sum_{i>\frac{n_2}{10t}}^{2tn_2+2t} A(i)$.

Let $f_1 \in T(f,n_2-n_1)$, $f_2 \in T(f,n_1)$, and let $j < n_3$.

By Lemma 11.1 the ideal generated by f^m is contained in the right ideal of $A \otimes A$, $\sum_{k=1}^{\infty}\sum_{j=0}^{\infty}((A(k\cdot n_3-n_2)\otimes A(j))T(f,n_2-n_1)T(f,n_1)(E(f,n_1,n_2)\cap E'(f,n_1,n_2,n_3))$. Let $f_1 \in T(f,n_2-n_1)$, $f_2 \in T(f,n_1)$, $g_3 \in E(f,n_1,n_2) \cap E'(f,n_1,n_2,n_3)$. By Lemma 9.3, there are $\frac{n_1-t}{2t} \leq \alpha_1 \leq 2tn_1+t$, $\frac{n_2-n_1-t}{2t} \leq \alpha_2 \leq 2t(n_2-n_1)+t$, such that $f_1 \in A(n_2-n_1)\otimes A(\alpha_1)$, $f_2 \in A(n_1)\otimes A(\alpha_2)$.

Assume first that $j+\alpha_1+\alpha_2 \in [kn_3-n_1, kn_3+n_1]$, for some k. Notice that the case $k=0$ is not possible as $j > a_2$ by Lemma 11.1. It follows that $(A(i\cdot n_3-n_2)\otimes A(j))f_1f_2g_3 \subseteq \sum_{j'\in U(n_1,n_3)}(A(n_3)\otimes A(j'))g_3(A\otimes A) \subseteq P$, because $g_3 \in E'(f,n_1,n_2)$.

If $j+\alpha_1+\alpha_2 \notin [kn_3-n_1, kn_3+n_1]$, for any k, it suffices to prove that $(A(n_3-n_2)\otimes A(j)f_1f_2T(f,n_2-n_1)T(f,a_1)(A(n_3)\otimes A(n_3)) \subseteq Q$ (notice that $A(n_3)$ at the end comes from the fact that f^m has large degree, and all considered sets are homogeneous).

Let $f_3 \in T(f,n_2-n_1)$, $f_4 \in T(f,n_1)$, then by Lemma 9.3, $f_3 \in A(n_1)\otimes A(\alpha_3)$, $f_4 \in A(n_2-n_1)\otimes A(\alpha_4)$, for some $\frac{n_1-t}{2t} \leq \alpha_3 \leq 2tn_1+t$, $\frac{n_2-n_1-t}{2t} \leq \alpha_4 \leq 2t(n_2-n_1)+t$. It suffices now to show that $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4(A(3tn_3)\otimes A(3tn_3)) \subseteq Q$.

It suffices now to consider the the following cases:

Case 1. $j+\alpha_1+\alpha_2 \in [kn_3+n_2+1, (k+1)n_3-n_2-1]$ for some $k \geq 0$. Then we have the following two subcases.

Case 1.a $j+\alpha_1+\alpha_2 \in [kn_3+n_2+1, (k+1)n_3-4tn_2]$ for some $k \geq 0$. Denote $i = j+\alpha_1+\alpha_2$, then $i+\alpha_3, \alpha_4 \leq (k+1)n_3-n_2-1$, since $\alpha_1+\alpha_2 \leq 2tn_2 = 2t$. Therefore, $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq \sum_{i:i,i+\alpha_3+\alpha_4\in[kn_3+n_2+1,(k+1)n_3-n_2-1]}(A(n_3)\otimes A(i))f_3f_4(A\otimes A) \subseteq Q(D,n_2,n_3) \subseteq Q$, since $D \subseteq F(f,n_1,n_2,n_3)$.

Case 1.b $j+\alpha_1+\alpha_2 \in [kn_3+4tn_2, (k+1)n_3-n_2-1]$ for some $k \geq 0$, then $j \geq kn_3+n_2+1$, since $\alpha_1+\alpha_2 \leq 2n_2+2t$. Hence, $((A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq \sum_{j:j,j+\alpha_1+\alpha_2\in[kn_3+n_2+1,(k+1)n_3-n_2-1]}(A(n_3)\otimes A(j))f_1f_2(A\otimes A) \subseteq Q(D,n_2,n_3) \subseteq Q$.

Case 2. $j+\alpha_1+\alpha_2 \in [kn_3+n_1+1, kn_3+n_2]$, for some k. We have the following two subcases.

Case 2a. $j+\alpha_1+\alpha_2 \in [kn_3+n_1+1, kn_3+n_2-4tn_1]$, for some k. In this case $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq \sum_{i\in[kn_3+n_1+1,kn_3+n_2-4tn_1]}(A(n_3)\otimes A(i))f_3(A\otimes A) \subseteq Q(T(f,n_1),n_1,n_2) \subseteq Q$, because $\alpha_3 \leq 2tn_1+t \leq 4tn_1-1$, and $f_3 \in T(f,n_1)$.

Case 2b. $j+\alpha_1+\alpha_2 \in [kn_3+n_2-4tn_1, kn_3+n_2]$, for some k. Denote $i=j+\alpha_1+\alpha_2$, then $i+\alpha_3+\alpha_4 \leq (k+1)n_3-n_2-1$.

By Lemma 12.1, we have $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq C$ where $C = \sum_{i:i,i+\alpha_3+\alpha_4\in[kn_3+n_2-4tn_1,(k+1)n_3-n_2-1],0\leq k'}(A(k'\cdot n_3)\otimes A(i))f_3f_4(A\otimes A)$.

Denote $q=\alpha_3+\alpha_4-5tn_1$, then $\bar{Z}_{5tn_1}(K\xi(f_3f_4)) \subseteq A(n_2)\otimes A(q)$. By Lemma 11.2, applied for $n=5tn_1$ and for the linear space Kf_3f_4, we get

$$C \subseteq \sum_{i:i,i+q\in[kn_3+n_2+tn_1-1,(k+1)n_3-n_2-1],0\leq k'} (A(k'\cdot n_3)\otimes A(i)))\bar{Z}_{5tn_1}(Kf_3f_4)(A\otimes A),$$

so $C \subseteq Q(F(f,n_1,n_2,n_3),n_2,n_3) \subseteq Q$, because $f_3f_4 \in D$, so $\bar{Z}_{5tn_1}(Kf_3f_4) \subseteq F(f,n_1,n_2,n_3)$.

Case 3. $j+\alpha_1+\alpha_2 \in [kn_3-n_2, kn_3-n_1-1]$, for some k. We have the following two subcases.

Case 3a. $j+\alpha_1+\alpha_2 \in [kn_3-n_2+4tn_1, kn_3-n_1-1]$, for some k.

Denote, $i=j+\alpha_1$, then $i+\alpha_2 = j+\alpha_1+\alpha_2 \leq kn_3-n_1-1$ and $i > (j+\alpha_1+\alpha_2)-\alpha_2 \geq (kn_3-n_2+4tn_1)-(2tn_1+2t) > kn_3-n_2+n_1+1=1$. Therefore the interval $[i,i+\alpha_2$ is disjoint with any interval $[k'n_2-n_1,k'n_2+n_1]$ for all k'. Hence, $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq \sum_{k=0}^{\infty}\sum_{i:i+\alpha_2\in[kn_3-n_2+n_1+1,kn_3-n_1-1]}(A(n_3-n_1)\otimes A(i))f_2(A\otimes A)$, hence $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq Q(T(f,n_1),n_1,n_2) \subseteq Q$.

Case 3b. $j+\alpha_1+\alpha_2 \in [kn_3-n_2, kn_3-n_2+4tn_1]$, for some k. Then, $j \geq kn_3-n_2-2tn_2-2t$, hence $j, j+\alpha_1+\alpha_2 \in [(k-1)n_3+n_2+1, kn_3-n_2+4tn_1]$. By Lemma 12.1, applied for the linear space Kf_1f_2, $p=5tn_1$ we have $(A(n_3-n_2)\otimes A(j))f_1f_2f_3f_4 \subseteq P_1$ where $P_1 = \sum_{j:j,j+\alpha_1+\alpha_2\in[(k-1)n_3+n_2+1,kn_3-n_2+4tn_1]}(A(n_3-n_2)\otimes A(i))f_1f_2(A(3tn_3)\otimes A(3tn_3))$. Denote $q=\alpha_1+\alpha_2-5tn_1$, then $Z_{5tn_1}(K\xi(f_1f_2)) \subseteq A(n_2)\otimes A(q)$. Observe that $P_1 \subseteq P_2$, where $P_2 = \sum_{j:j,j+q\in[(k-1)n_3+n_2+1,kn_3-n_2-1]}(A(n_3-n_2)\otimes A(i))Z_{5tn_1}(Kf_1f_2)(A(3tn_3)\otimes A(3tn_3))$. Notice that $f_1f_2 \in D$, and so $Z_{5tn_1}(Kf_1f_2) \subseteq F(f,n_1,n_2,n_3)$. By Lemma 12.1, $f_1f_2 \subseteq A(n_2)\otimes A$, hence $P_2 \subseteq Q(F(f,n_1,n_2),n_2,n_3) \subseteq Q$. □

Let $1 \leq \alpha$, $0 < n_1 < n_2$ be natural numbers and let $F \subseteq A(\alpha)\otimes A(n_1)$. Recall that

$$Q'(F,n_1,n_2) = \sum_{k=0}^{\infty}\sum_{j\in W(n_1,n_2)}(A(j)\otimes(A(k\cdot n_2)+A(k\cdot n_2-n_1))F(A\otimes A),$$

where $j \in W(n_1,n_2)$ if and only if $[j,j+\alpha]$ is disjoint with all the intervals $[kn_2-n_1,kn_2+n_1]$ for $k=0,1,2,\ldots$.

Let $x_i \in \{x,y\}$ be generators of A, denote $(x_1x_2\cdots x_n)^{op} = x_nx_{n-1}\cdots x_1$. Let $a_i,b_i \in A$. For $r=\sum_i a_i\otimes b_i$ define $\xi(r)=\sum_i a_i\otimes(b_i)^{op}$. The following Lemma is similar to Lemma 11.3

Recall that if $a \in A(n) \otimes A(m)$, for some m, n, then $l(a) = n$, $r(a) = m$. We are now ready to prove the two main theorems in this section.

Lemma 11.4 *Let $f \in H$ where $H = \sum_{1 \leq i,j \leq t} A(i) \otimes A(j)$ for some $t \geq 1$. Let n_1, n_2, n_3 be natural numbers such that n_1 divides n_2, n_2 divides n_3 and $10tn_1 < n_2$, $10tn_2 < n_3$, $10tn_3 < m$, $10t < n_1$. Let $U(n_1, n_3) = \bigcup_{k=1}^{\infty}[kn_3 - n_1, kn_3 + n_1]$. Let $G_1(f, n)$, $G_2(f, n)$ be as in Lemma 10.2, and $T(f, n)$, $T'(f, n)$ be as in Lemmas 9.3 and 9.5 and $E'(f, n_1, n_2)$ be as in Lemma 11.1 (and let $\xi(r)$ be defined as above for any $r \in A \otimes A$.) Then there is set $F'(f, n_1, n_2, n_3) \subseteq A(n_2) \otimes \sum_{i=1}^{2tn_2+2t} A(i)$ with $\dim_K F(f, n_1, n_2, n_3) \leq 2^{40tn_1} n_2$ and such that the two-sided ideal of $A \otimes A$ generated by $\xi(f^m)$ belongs to the right ideal of $A \otimes A$ generated by elements from $Q' + P'$ where $Q' = Q(\xi(T(f, n_1)), n_1, n_2)) + Q(F'(f, n_1, n_2, n_3), n_2, n_3)$, and*

$$P' = \sum_{\alpha=0}^{\infty} \sum_{k=1}^{\infty} \sum_{j+\alpha \in U(n_1,n_3)} (A(k \cdot n_3) \otimes A(j)) \xi(E'(f, n_1, n_2, \alpha))(A \otimes A),$$

where $E'(f, n_1, n_2) = \sum_{i=0}^{\infty} E'(f, n_1, n_2, \alpha)$, with $E'(f, n_1, n_2, \alpha) \subseteq A \otimes A(\alpha)$. Moreover, if $n_2 > 100tn_1$ then $F(f, n_1, n_2, n_3) \subseteq A(n_2) \otimes \sum_{i > \frac{n_2}{10t}}^{2tn_2+2t} A(i)$.

Proof. Let $D = \xi(T(f, n_2 - n_1)T(f, n_1) + T(f, n_1)T(f, n_2 - n_1) + T(f, n_2))$,

$$F'(f, n_1, n_2, n_3) = D + Z_{5tn_1}(D) + \bar{Z}_{5tn_1}(D),$$

where $Z_n(D)$, $\bar{Z}_n(D)$ are as in Lemma 11.2. By Lemma 11.2, we get $\dim_K F'(f, n_1, n_2, n_3) \leq (1 + 2^{5tn_1+1}) \dim_K D \leq 2^{40tn_1} n_2$. By Lemma 9.3, $T(f, n), \xi(T(f, n)) \subseteq A(n) \otimes \sum_{i > \frac{n-t}{2t}}^{2nt+t} A(i)$, hence $F'(f, n_1, n_2, n_3) \subseteq A(n_2) \otimes \sum_{i > \frac{n_2-2t}{2t} - 5tn_1}^{2tn_2+2t} A(i)$. Notice that if $n_2 > 100tn_1$ then $F'(f, n_1, n_2, n_3) \subseteq A(n_2) \otimes \sum_{i > \frac{n_2}{10t}}^{2tn_2+2t} A(i)$.

Let $f_1 \in T(f, n_2 - n_1)$, $f_2 \in T(f, n_1)$, and let $j < n_3$. By Lemma 9.3, $f_1 \in A(n_2 - n_1) \otimes A(\alpha_1)$, $f_2 \in A(n_1) \otimes A(\alpha_2)$, for some $\alpha_1 \leq 2tn_1 + t$, $\frac{n_2 - n_1 - t}{2t} \leq \alpha_2 \leq 2t(n_2 - n_1) + t$.

Let J denote the ideal generated by $\xi(f^m)$ in $A \otimes A$, and let J' denotes the ideal generated by f^m in $A \otimes A$. By Lemma 11.1, $\xi(J') \subseteq \sum_{k,j=0}^{\infty} \xi((A(k \cdot n_3 - n_2) \otimes A(j))T(f, n_2 - n_1)T(f, n_1)(E(f, n_1, n_2) \cap E'(f, n_1, n_2, n_3)))(A \otimes A)$. Hence, $J \subseteq \sum_{k,j'=0}^{\infty}((A(k \cdot n_3 - n_2) \otimes A(j'))\xi(T(f, n_2 - n_1)T(f, n_1)(E(f, n_1, n_2) \cap E'(f, n_1, n_2, n_3)))(A \otimes A)$. Recal that sets $E(f, n_1, n_2)$ and $E'(f, n_1, n_2)$ are homogeneous, and hence they are generated by homogeneous elements.

Let $f \in A(s) \otimes A(j)$ be such that $f \in E(f, n_1, n_2) \cap E'(f, n_1, n_2, n_3)$. Assume first that $j + j' \in [k'n_3 - n_1, k'n_3 + n_1]$, for some k'. Notice that

the case $k' = 0$ is not possible as elements from $E(f, n_1, n_2)$ have right degree larger than n_1. It follows that $(A(k \cdot n_3 - n_2) \otimes A(j'))\xi(f_1 f_2 f) \subseteq (A(k \cdot n_3) \otimes A(j')\xi(f)(A \otimes A) \subseteq P'$, because $\xi(f) \in \xi(E'(f, n_1, n_2, j))$ and $j + j' \in U(n_1, n_3)$.

Assume now that $j + j' \notin [k'n_3 - n_1, k'n_3 + n_1]$, for any k'. Recall that $f \in E(f, n_1, n_2)$, so $f = f_3 f_4 f_5$ for some $f_3 \in A(n_1) \otimes A(\alpha_3)$, $f_4 \in A(n_2 - n_1) \otimes A(\alpha_4)$, $f_5 \in A(\beta) \otimes A(\alpha_5)$ for some $\alpha_3 \leq 2tn_1 + t$, $\frac{n_2 - n_1 - t}{2t} \leq \alpha_4 \leq 2t(n_2 - n_1) + t$, and some β, α_5. Observe that $\alpha_3 + \alpha_4 + \alpha_5 = j$ because $f_3 f_4 f_5 = f$.

Notice that it suffices to show that $(A(n_3 - n_2) \otimes A(j'))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq Q$. We have the following cases:

Case 1. $j + j' \in [kn_3 + n_2 + 1, (k+1)n_3 - n_2]$ for some $k \geq 0$. Then we have the following two subcases.

Case 1.a $j + j' \in [kn_3 + n_2 + 1, (k+1)n_3 - 4tn_2]$ for some $k \geq 0$. Then, $(A(n_3 - n_2) \otimes A(j'))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq \sum_{kn_3 + n_2 < j + j', j + j' + \alpha_1 + \alpha_2 < (k+1)n_3 - n_2} (A(n_3 - n_2) \otimes A(j + j'))\xi(f_1 f_2)(A \otimes A) \subseteq Q(D, n_2, n_3) \subseteq Q$, since $\xi(f_1 f_2) \in D \subseteq F(f, n_1, n_2, n_3)$.

Case 1.b $j + j' \in [kn_3 + 4tn_2, (k+1)n_3 - n_2]$ for some $k \geq 0$. We have, $((A(n_3 - n_2) \otimes A(j))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq \sum_{kn_3 + 2n_2 < j + j', j + j' - \alpha_3 - \alpha_4) < (k+1)n_3 - n_2} (A(n_3) \otimes A(j + j' - \alpha_3 - \alpha_4)\xi(f_3 f_4)(A \otimes A) \subseteq Q(D, n_2, n_3) \subseteq Q$, since $\xi(f_3 f_4) \in D$.

Case 2. $j + j' \in [kn_3 + n_1 + 1, kn_3 + n_2]$, for some k. We have the following two subcases.

Case 2a. $j + j' \in [kn_3 + n_1 + 1, kn_3 + n_2 - 4tn_1]$, for some k. In this case $(A(n_3 - n_2) \otimes A(j))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq \sum_{j, j' : j + j' \in [kn_3 + n_1 + 1, kn_3 + n_2 - 4tn_1]} (A(n_3 - n_1) \otimes A(i))\xi(f_2)(A \otimes A) \subseteq Q(\xi(T(f, n_1)), n_1, n_2) \subseteq Q$ (because $\alpha_3 \leq 2tn_1 + t \leq 4tn_1 - n_1 - 1$, so $j + j', j + j' + \alpha_3 \in [kn_3 + n_1 + 1, kn_3 + n_2 - n_1 - 1]$).

Case 2b. $j + j' \in [kn_3 + n_2 - 4tn_1, kn_3 + n_2]$, for some k. Denote $i = j + j'$. By Lemma 12.1, we have $(A(n_3 - n_2) \otimes A(j))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq E$ where $E = \sum_{i : i, i + \alpha_1 + \alpha_2 \in [kn_3 + n_2 - 4tn_1, (k+1)n_3 - n_2 - 1], 0 \leq k'} (A(k' \cdot n_3 - n_2) \otimes A(i))\xi(f_1 f_2)(A \otimes A)$.

Let $q = \alpha_1 + \alpha_2 - 5tn_1$, then $\bar{Z}_{5tn_1}(K\xi(f_1 f_2)) \subseteq A(n_2) \otimes A(q)$. By Lemma 11.2, applied for $n = 5tn_1$ and for the linear space $K\xi(f_1 f_2)$, we get

$$E \subseteq \sum_{i : i, i + q \in [kn_3 + n_2 + 1, (k+1)n_3 - n_2 - 1], 0 \leq k'} (A(k' \cdot n_3) \otimes A(i)))\bar{Z}_{5tn_1}(K\xi(f_1 f_2))(A \otimes A).$$

Observe now that $\xi(f_1 f_2) \in D$, so $\bar{Z}_{5tn_1}(K\xi(f_1 f_2)) \subseteq F'(f, n_1, n_2, n_3)$ hence $E \subseteq Q(F'(f, n_1, n_2, n_3), n_2, n_3) \subseteq Q$,

Case 3. $j + j' \in [kn_3 - n_2, kn_3 - n_1 - 1]$, for some k. We have the following two subcases.

Case 3a. $j + j' \in [kn_3 - n_2 + 4tn_1, kn_3 - n_1 - 1]$, for some k. Denote $i = j + j'$. Then $[i - \alpha_3, i]$ is disjoint with any interval $[k'n_2 - n_1, k'n_2 + n_1]$, because

$i \leq kn_3 - n_1 - 1$, and $i - \alpha_3 \geq kn_3 - n_2 + 4tn_1 - \alpha_3 > kn_3 - n_2 + n_1 + 1$. Hence, $(A(n_3 - n_2) \otimes A(j))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq \sum_k \sum_{i:i, i-\alpha_3 \in [kn_3 - n_2 + n_1 + 1, kn_3 - n_1 - 1]} (A(n_3) \otimes A(i - \alpha_3))\xi(f_3)(A \otimes A) \subseteq Q(\xi(T(f, n_1)), n_1, n_2) \subseteq Q$.

Case 3b. $j + j' \in [kn_3 - n_2, kn_3 - n_2 + 4tn_1]$, for some k. Denote $i = j + j' - \alpha_3 - \alpha_4$. Then, $i, i + \alpha_3 + \alpha_4 \in [kn_3 - n_2 - 2tn_2 - 2t, kn_3 - n_2 + 4tn_1]$, and so $i, i + \alpha_3 + \alpha_4 \in [(k-1)n_3 + n_2 + 1, kn_3 - n_2 + 4tn_1]$. By Lemma 12.1, applied for the linear space $K\xi(f_3 f_4)$, $p = 5tn_1$ we have $(A(n_3 - n_2) \otimes A(j))\xi(f_1 f_2 f_3 f_4 f_5) \subseteq P_1$ where $P_1 = \sum_{i:i, i+\alpha_3+\alpha_4 \in [(k-1)n_3 + n_2 + 1, kn_3 - n_2 + 4tn_1]} (A(n_3) \otimes A(i))\xi(f_3 f_4)(A \otimes A)$. Denote $q = \alpha_3 + \alpha_4 - 5tn_1$, then $Z_{5tn_1}(K\xi(f_3 f_4)) \subseteq A(n_2) \otimes A(q)$. It follows that $P_1 \subseteq P_2$ where

$$P_2 = \sum_{i:i, i+q \in [(k-1)n_3 + n_2 + 1, , kn_3 - n_2 - 1]} (A(n_3) \otimes A(i)) Z_{5tn_1}(K f_1 f_2)(A \otimes A).$$

Notice that $\xi(f_3 f_4) \in D$, and so $Z_{5tn_1}(K\xi(f_3 f_4)) \subseteq F'(f, n_1, n_2, n_3)$, hence $P_2 \subseteq Q(F(f, n_1, n_2), n_2, n_3) \subseteq Q$. □

12 Appendix: Proof of Theorem 3.1. Constructing space S

In this section we will show that there is a nil ideal N in $A \otimes A$ such that $N \subseteq S$ for some 'good' linear space $S \subseteq A \otimes A$. Space S will be constructed in Corollary 3.5 in this section.

We will start with a supporting lemma which can be read independently of the rest of the paper.

Lemma 12.1 *Let $n_2, i, i', \alpha, \beta$ be natural numbers such that $4p < n_2$, $-p \leq i, i' \leq p$, $0 \leq \alpha, \beta \leq p$. Let $G \subseteq A(n_2 - \alpha) \otimes A(n_2 - \beta)$ be a linear space. Then there is set $Z_{i,i'}(G) \subseteq A(n_2) \otimes A(n_2)$ with $\dim_K Z_{i,i'}(G) < 2^{6p+2} \dim_K G$ and such that for all $k, k' \geq p$,*

$$(A(k + i) \otimes A(k' + i'))G(A(3p) \otimes A(3p)) \subseteq (A(k) \otimes A(k'))Z_{i,i'}(G)(A \otimes A).$$

Proof. We have four cases: **Case** 1. The case when $i, i' > 0$. Denote $E = (A(i) \otimes A(i'))G(A(\alpha) \otimes A(\beta))$. Then $\dim_K E \leq 2^{4p} \dim_K G$. Note that $E \subseteq A(n_2 + i) \otimes A(n_2 + i')$. Similarly, as in Lemma 10.1 there is set $T(E) \subseteq A(n_2) \otimes A(n_2)$ such that $E \subseteq T(E)(A(i) \otimes A(i'))$ and $\dim_K(T'(E)) \leq 2^{2p+2} \dim_K E \leq 2^{6p+2} \dim_K G$. We will set then $Z_{i,i'} = T(E)$. Observe that by construction, for every $j, j' \geq 0$, we have $(A(k + i) \otimes A(k' + i'))G(A(3p) \otimes A(3p)) \subseteq (A(k) \otimes A(k'))E(A(3p - \alpha - i) \otimes A(3p - \beta - i')) \subseteq (A(k) \otimes A(k'))Z_{i,i'}(A \otimes A)$, as required.

Case 2. The case when $i, i' \leq 0$. Denote $E = \sum_{\alpha,\beta} G(A(-i+\alpha) \otimes A(-i'+\beta)$. Then $\dim_K E \leq 2^{4p} \dim_K G$. Note that $E \subseteq A(a_2-i) \otimes A(a_2-i')$ (where $i, i' \leq 0$). By an analogon of Lemma 10.1 there is set $T'(E) \subseteq A(a_2) \otimes A(a_2)$ such that $E \subseteq (A(-i) \otimes A(-i'))T'(E)$ and $\dim_K(T'(E)) \leq 2^{2p+2} \dim_K E \leq 2^{6p+2} \dim_K G$. We will set then $Z_{i,i'} = T'(E)$. Observe that by construction, for every $j, j' \geq 0$,

$$A(k+i) \otimes A(k'+i'))G(A(3p) \otimes A(3p)) \subseteq (A(k) \otimes A(k'))Z_{i,i'}(A \otimes A),$$

as required.

Case 3. The case when $i > 0$, $i' \leq 0$. Denote $E = (A(i) \otimes K)G(A(\alpha) \otimes A(\beta - i')$. Then $\dim_K E \leq 2^{4p} \dim_K G$. Note that $E \subseteq A(a_2+i) \otimes A(a_2-i')$ (where $i' \leq 0$). By a similar argument as in Lemma 10.1 there is set $T''(E) \subseteq A(a_2) \otimes A(a_2)$ such that $E \subseteq (K \otimes A(-i'))T''(E)(A(i) \otimes K)$ and $\dim_K(T''(E)) \leq 2^{2p+2} \dim_K E \leq 2^{6p+2} \dim_K G$. We will set then $Z_{i,i'} = T''(E)$. Observe that by construction, for every $j, j' \geq 0$,

$$(A(k+i) \otimes A(k'+i'))G(A(3p) \otimes A(3p)) \subseteq (A(k) \otimes A(k'))Z_{i,i'}(A \otimes A),$$

as required.

Case 4. The case when $i \leq 0$, $i' > 0$. This case is done similarly as case 3. □

We are now ready to prove the following theorem.

Theorem 12.1 *Let $f \in \sum_{0<i,j\leq t} A(i) \otimes A(j)$ for some $t \geq 1$. Let n_1, n_2, n_3, m be natural numbers such that n_1 divides n_2, n_2 divides n_3, $10tn_1 < n_2$, $10tn_2 + 2n_3 < n_3$, $10tn_3 + 2n_3 < m$, $10t < n_1$. Let $T(f, n_1)$, $E'(f, n_1, n_2)$, $F(f, n_1, n_2, n_3)$, $F'(f, n_1, n_2, n_3)$ be as in Lemmas 11.3, 9.6 and 11.4. Then there exists a homogeneous linear space $G(f, n_1, n_2) \subseteq A(n_2) \otimes A(n_2)$ with $\dim_K G(f, n_1, n_2) < n_2^3 2^{500tn_1}$ and such that, for every integer $m > 20n_3$, the ideal generated by f^m and $\xi(f^m)$ in $A \otimes A$ is contained in the right ideal of $A \otimes A$, $M + Q_1 + Q_2 + Q_3$ where $Q_1 = Q(T(f, n_1) + \xi(T(f, n_1)), n_1, n_2)$, $Q_2 = Q(F(f, n_1, n_2, n_3), n_2, n_3) + Q'(F'(f, n_1, n_2, n_3), n_2, n_3)$, $Q_3 = Q'(T'(f, n_1) + \xi(T'(f, n_1)), n_1, n_2)$ and where $M = \sum_{i=1,2,\ldots;j=1,2,\ldots}((A(i \cdot n_3) + A(i \cdot n_3 - n_2)) \otimes (A(j \cdot n_3) + A(j \cdot n_3 - n_2)))G(f, n_1, n_2)(A \otimes A)$.*

Proof. Let I' be the ideal generated in $A \otimes A$ by $\xi(f^{m-2n_3})$ and f^{m-2n_3}, and I be the ideal generated by $\xi(f^m)$ and f^m. Then $I \subseteq I'(A(n_3) \otimes A(n_3))$, since $\xi(f^m) = \xi(f^{n_3} \cdot \xi(f^{m-2n_3}) \cdot f^{n_3})$. So we only need to show that $I'(A(n_3) \otimes A(n_3)) \subseteq M + Q_1 + Q_2 + Q_3$.

By Lemmas 11.3 and 11.4, the two sided ideal generated by f^{m-2n_3} and $\xi(f^{m-2n_3})$ belongs to the right ideal of $A \otimes A$ generated by elements

from Q_1, Q_2 and from $P+P'$ where $P=\sum_{k=1}^{\infty}\sum_{j'\in U(n_1,n_3)}(A(k\cdot n_3)\otimes A(j'))(E'(f,n_1,n_2)$, and from $P'=\sum_{\alpha=0}^{\infty}\sum_{k=1}^{\infty}\sum_{j':j'+\alpha\in U(n_1,n_3)}(A(k\cdot n_3)\otimes A(j))\xi(E'(f,n_1,n_2,\alpha))$. So we only need to show that $(P+P')(A(n_3)\otimes A(n_3))\subseteq M+Q_1+Q_2+Q_3$.

Let $G_1(f,n_2-n_1)$, $G_2(f,n_2-n_1)$ be as in Theorem 10.2 applied for $n=n_2-n_1$, then we can write $G_1(f,n_2-n_1)=\sum_{\gamma\geq\frac{n_2-n_1}{2t}}^{n_2-n_1}G_\gamma$ for some $G_i\subseteq A(n_2-n_1)\otimes A(i)$. Let c be the smallest number such that

$$c\geq\frac{n_2-n_1}{2t}.$$

Write now $G_1(f,n_2-n_1)=G+G'$ where $G=\sum_{i=c}^{n_2-10tn_1}G_i$, $G'=\sum_{i=n_2-10tn_1+1}^{n_2-n_1}G_i$. Similarly, write $G_2(f,n_2-n_1)=\bar{G}+\bar{G}'$ where $\bar{G}=\sum_{i=c}^{n_2-10tn_1}\bar{G}_i$, $\bar{G}'=\sum_{i=n_2-10t_1+1}^{n_2-n_1}\bar{G}_i$, for suitable $\bar{G}_i\subseteq A(i)\otimes A(n_2-n_1)$.

Define now $F_1(f,n_1,n_2)=\sum_{j=n_2-10tn_1}^{n_2-n_1}(D_j+\bar{D}_j)$ where

$$D_j=\sum_{-p\leq i,i'\leq p}Z_{i,i'}(G_j),\bar{D}_j=\sum_{-p\leq i,i'\leq p}Z_{i,i'}(\bar{G}_j),$$

where $Z_{i,i'}(G_j)$ is as in Lemma 12.1 applied for $p=12tn_1$, for the same n_2, and for $G=G_j$ (and $Z_{i,i'}(\bar{G}_j)$ for $G=\bar{G}_j$). By Lemma 12.1, we have $F_1(f,n_1,n_2)\subseteq A(n_2)\otimes A(n_2)$ and $\dim_K F_1(f,n_1,n_2)<n_2^3 2^{500tn_1-1}$.

Denote $L_{j'}=\sum_{j\in U(n_1,n_3)}(A(j'n_3)\otimes A(j))(G+\bar{G})(A(n_3)\otimes A(n_3))$. Denote $p=12tn_1$. Observe now that if $j\in U(n_1,n_3)$ then there exists k',i' such that $-p\leq i'\leq p$, k' is divisible by n_3 and $j=k'+i'$. Apply Lemma 12.1 for this k',i' and p, and for for $k=j'n_3$, $i=0$. We then get for every j', $L_{j'}\subseteq(A(k)\otimes A(k'))G_1(f,n_1,n_2)(A\otimes A)\subseteq M$, provided that $F_1(f,n_1,n_2)\subseteq G(f,n_1,n_2)$.

Similarly, define $F_2(f,n_1,n_2)=\sum_{j=n_2-(t+1)n_1}^{n_2-n_1}(D'_j+\bar{D}'_j)$ where

$$D'_j=\sum_{-p\leq i,i'\leq p}Z_{i,i'}(\xi(G_j)),\bar{D}'_j=\sum_{-p\leq i,i'\leq p}Z_{i,i'}(\xi(\bar{G}_j)),$$

where $Z_{i,i'}(\xi(G_j))$ is as in Lemma 12.1 applied for $p=12tn_1$, for the same n_2,n_3, and for $G=\xi(G_j)$ (and $Z_{i,i'}(\xi(\bar{G}_j))$ for $G=\xi(\bar{G}_j)$). Similarly as before, $F_2(f,n_1,n_2)\subseteq A(n_2)\otimes A(n_2)$ and $\dim_K F_2(f,n_1,n_2)\leq n_2^3 2^{500tn_1-1}$. For $g\in\xi(G)+\xi(\bar{G})$ denote $\bar{L}_{j',g}=\sum_{j:j+r(g)\in U(n_1,n_3)}(A(j'n_3)\otimes A(j))g(A(n_3)\otimes A(n_3))$. Denote $p=12tn_1$. Observe now that since $j+r(g)\in U(n_1,n_2)$ and $g\in\sum_{l,l'=n_2-10tn_1}^{n_2-n_1}A(l)\otimes A(l')$ then there exists k' such that $k'+n_2$ is divisible by n_3, and $-p\leq i'\leq p$, such that $j=k'+i'$. Apply Lemma 12.1 for this k',i' and p, and for for $k=j'n_3$, $i=0$. We then get for

every j', g', $L_{j',g} \subseteq (A(k) \otimes A(k'))F_2(f, n_1, n_2)(A \otimes A) \subseteq M$, provided that $F_2(f, n_1, n_2) \subseteq G(f, n_1, n_2)$. Let now h be a homogeneous element such that $h \in g'(A(l) \otimes A(l'))$ for some $g' \in G + \bar{G}$ and for some l, l', and denote $g = \xi(g')$. Observe that then $\xi(h) \in (K \otimes A(l')g(A(l) \otimes K)$. Therefore, $\sum_{j:j+r(h)\in U(n_1,n_3)}(A(j'n_3) \otimes A(j))\xi(h)(A(n_3) \otimes A(n_3)) \subseteq \sum_{j:j+l'+r(g)\in U(n_1,n_3)}(A(j'n_3) \otimes A(j + l'))\xi(g)(A(n_3) \otimes A(n_3))(A \otimes A) \subseteq M$, for some k'' divisible by a_{n_3}, provided that $F_2(f, n_1, n_2) \subseteq G(f, n_1, n_2)$, as before (with $j + l'$ instead of j).

We now set $G(f, n_1, n_2) = F_1(f, n_1, n_2) + F_2(f, n_1, n_2)$.

Recall that $E'(f, n_1, n_2) = S(f, n_1, n_2)+S'(f, n_1, n_2)$, as in Lemma 11.1.

We will now show that $(P + P')(A(n_3) \otimes A(n_3) \subseteq M + Q_1 + Q_2 + Q_3$. by considering all the possible cases :

Case 1. To show that $\sum_{j'=1}^{\infty}\sum_{j'\in U(n_1,n_3)}(A(j'\cdot n_3)\otimes A(j))S(f, n_1, n_2)(A(n_3)\otimes A(n_3)) \subseteq M + Q_1 + Q_2$. Notice that $S(f, n_1, n_2) = G_1(f, n_2 - n_1)T(f, n_1) = GT(f, n_1) + G'T(f, n_1)$, so we consider the following subcases

Case 1a To prove that for all $j' > 0$, we have $\sum_{j\in U(n_1,n_3)}(A(j'n_3 \otimes A(j))GT(f, n_1)(A(n_3) \otimes A(n_3)) \subseteq M$, this follows from the first part of this proof.

Case 1b To prove that for all $j' > 0$, we have $\sum_{j\in U(n_1,n_3)}(A(j'n_3 \otimes A(j))G'T(f, n_1) \subseteq Q_1$. By assumption $c \geq \frac{n_2-n_1}{2t} > 2n_1$. Observe that since $c \leq \beta \leq n_2 - 10tn_1$, and $j \in U(n_1, n_3)$ then interval $[j + \beta, j + \beta + n_1]$ is disjoint with any interval $[kn_2 - n_1, kn_2 + n_1]$. It follows that then $\sum_{j\in U(n_1,n_3)}(A(j'n_3 \otimes A(j))G_\beta T(f, n_1) \subseteq \sum_{j\in U(n_1,n_3)}(A(j'n_3 + n_2 - n_1) \otimes A(j + \beta))T(f, n_1) \subseteq Q(T(f, n_1), n_1, n_2) \subseteq Q_1$.

Case 2 To show that for all $j' > 0$, $\sum_{1\leq j',\alpha}\sum_{j+\alpha\in U(n_1,n_3)}(A(j' \cdot n_3) \otimes A(j))(\xi(S(f, n_1, n_2) \cap (A \otimes A(\alpha))(A(n_3) \otimes A(n_3)) \subseteq M + Q_1 + Q_2$. Notice that $\xi(S(f, n_1, n_2)) = \xi(GT(f, n_1)+G'T(f, n_1))$, so we consider the following subcases

Case 2a To show that $L_{j'} \subseteq M$ for all $j' > 0$, and all homogeneous $h \in \xi(GT(f, n_1)(A \otimes A))$ where

$$L_{j',h} = \sum_{j+r(h)\in U(n_1,n_3)} (A(j' \cdot n_3) \otimes A(j))\xi(h)(A(n_3) \otimes A(n_3)).$$

Observe that $L_{j',h} \subseteq M$ by the first part of this proof.

Case 2b To prove that $L_{j',g} \subseteq Q_1$ for every every $j' > 0$, and every homogeneous $g \in \xi(GT(f, n_1)(A\otimes A))$ where $L_{j',g} = \sum_{j:j+r(g)\in U(n_1,n_3)}(A(j'n_3)\otimes A(j))g$. Recall that $r(g)$ is the right degree of g, so $r(g) = i$ if $g \in A \otimes A(i)$. Observe that if $g = \xi(g_1g_2g_3)$ where $g_1 \in G_\beta, g_2 \in T(f, n_1)$, $g_3 \in A \otimes A$ are homogeneous elements, then $L_{j',g} \subseteq \sum_{l:l+\beta+r(g_2)\in U(n_1,n_3)}(A(j'n_3 + n_2 - n_1) \otimes A(l))\xi(g_2)(A \otimes A)$, (where $l = j + r(g_3)$).

By assumption $2n_1 < c \leq \frac{n_2-n_1}{2t}$ and $c \leq \beta \leq n_2 - 10tn_1$. Recall also that $r(g_2) \leq 2tn_1 + t$, by Lemma 9.3. Therefore $l + \beta + r(g_2) \in U(n_1, n_3)$ yields that the interval $[l, l+r(g_2)]$ is disjoint with any interval $[kn_2 - n_1, kn_2 + n_1]$. It follows that $L_{j',g} \subseteq Q(\xi(T(f, n_1)), n_1, n_2) \subseteq Q_1$.

Case 3 To show that for all $j' > 0$, $\sum_{j'=0}^{\infty} \sum_{j \in U(n_1,n_3)} (A(j' \cdot n_3) \otimes A(j)) S'(f, n_1, n_2)(A(n_3) \otimes A(n_3)) \subseteq M + Q_1 + Q_2 + Q_3$.

By Lemma 11.1, $S'(f, n_1, n_2) = G_2(f, n_2 - n_1)S = \bar{G}S + \bar{G}'S$, where $S = \bigcap_{i=1}^{2n_1} T'(f, i)T(f, n_1)(A \otimes A)$, so we consider the following subcases

Case 3a To show that for all $j' > 0$, $\sum_{j'=0}^{\infty} \sum_{j \in U(n_1,n_3)} (A(j' \cdot n_3) \otimes A(j)) \bar{G}S(A(n_3) \otimes A(n_3)) \subseteq M$. This follows from the first part of this proof where we showed that $\sum_{j \in U(n_1,n_3)} (A(j'n_3) \otimes A(j))(G + \bar{G})(A(n_3) \otimes A(n_3)) \subseteq M$.

Case 3b To show that for all $j' > 0$, $L_{j'} \subseteq Q_3$, where $L_{j'} = \sum_{j \in U(n_1,n_3)} (A(j' \cdot n_3) \otimes A(j)) \bar{G}'S(A(n_3) \otimes A(n_3))$. Let k' be divisible by n_3 and such that $-n_1 \leq k' - j \leq n_1$ (because $j \in U(n_1, n_2)$ such k' exists). Notation $k' \in U$ will mean that k' is divisible by n_3.

Denote $\alpha = n_1 + k' - j$, then $\alpha + (n_2 - n_1 + j) = k' + n_2$, and $0 \leq \alpha \leq 2n_1$. We get $L_{j'} \subseteq \sum_{\beta=c}^{n_2-10tn_1} J_\beta$ where $J_\beta = ((A(j' \cdot n_3) \otimes A(j))\bar{G}_\beta T'(f, \alpha)T'(f, n_1)(A \otimes A)$. Recall that $T'(f, \alpha) \subseteq \sum_{i=1}^{2t\alpha+t} A(i) \otimes A(n_1)$, (by Lemmas 9.5, 9.6) hence

$$J_\beta \subseteq \sum_{i=1}^{2t\alpha+t} \sum_{k' \in U} (A(j' \cdot n_3 + \beta + i) \otimes A(k' + n_2))T'(f, n_1)(A \otimes A)$$

(because $k' + n_2 = j + (n_2 - n_1) + \alpha$). Recall that $2n_1 < c \leq \beta \leq n_2 - 10tn_1, \alpha < 2n_1$, $T'(f, n_1) \subseteq \sum_{i'=1}^{2tn_1+t} A(n_1) \otimes A(i')$. Observe that for $i \leq 4tn_1 + t, i' \leq 2tn_1 + t$, the interval $[j' \cdot n_3 + \beta + i, j' \cdot n_3 + \beta + i + i']$ is disjoint with any interval $[kn_2 - n_1, kn_2 + n_1]$. It follows that $J_\beta \subseteq Q'(T'(f, n_1), n_1, n_2) \subseteq Q_3$.

Case 4 To show that for all $j' > 0$, $L_{j'} \subseteq Q_3$ for every homogeneous $g \in \xi(S'(f, n_1, n_2))$ where $L_{j',g} = \sum_{j:j+r(g) \in U(n_1,n_3)} (A(j'n_3) \otimes A(j))g(A(n_3) \otimes A(n_3))$. (recall that $r(g)$ is the right degree of g).

By Lemma 11.1, $S'(f, n_1, n_2) = G_2(f, n_2 - n_1)S = \bar{G}S + \bar{G}'S$, where $S = \bigcap_{i=0}^{2n_1} T'(f, i)T(f, n_1)(A \otimes A)$. So we consider the following subcases

Case 4a To show that for all $j' > 0$, and any $h \in \bar{G}S$ we have $L_{j',h} \subseteq M$ where $L_{j',h} = \sum_{j:j+r(h) \in U(n_1,n_3)} (A(j' \cdot n_3) \otimes A(j))\xi(h)(A(n_3) \otimes A(n_3)) \subseteq M$. This follows from the first part of this proof.

Case 4b To prove that $L_{j',g} \subseteq Q_3$ for homogeneous $g \in \bar{G}'S$ and every $j' > 0$, where $L_{j',g} = \sum_{j:j+r(g) \in U(n_1,n_3)} (A(j'n_3) \otimes A(j))\xi(g)$. Since $j + r(g) \in U(n_1, n_3)$, then there is k' is divisible by n_3 and such that $-n_1 \leq j + r(g) - k' \leq n_1$. Denote $\alpha = n_1 + j + r(g) - k'$, then $0 \leq \alpha \leq 2n_1$.

Since $g \in \bar{G}'S$ then $g \in \bar{G}_\beta T'(f,\alpha)T(f,n_1)(A\otimes A)$, for some $c \le \beta \le n_2 - 10tn_1$. Therefore $g = g_1g_2g_3g_4$ where $g_1 \in \bar{G}_\beta$, $g_2 \in T'(f,\alpha)$, $g_3 \in T'(f,n_1)$, $g_4 \in A\otimes A$ are homogeneous elements. It follows that $L_{j',g} \subseteq \sum_{j:j+(r(g_1)+r(g_2)+r(g_3)+r(g_4))\in U(n_1,n_3)}((A(j'\cdot n_3+\beta+l(g_2))\otimes A(j+r(g_4))\xi(g_3)(A\otimes A)$. Notice that $r(g_1)+r(g_2) = n_2-n_1+\alpha$.

Recall that $g_2 \in T'(f,\alpha) \subseteq \sum_{i=1}^{2t\alpha+t} A(i)\otimes A(\alpha)$. Consequently, $L_{j',g} \subseteq \sum_{i=1}^{2t\alpha+t}\sum_{j:j+n_2-n_1+\alpha+r(g_3)+r(g_4))\in U(n_1,n_3)}((A(j'\cdot n_3+\beta+i)\otimes A(j+r(g_4))\xi(g_3)(A\otimes A)$. Observe that $j+r(g_3)+r(g_4) = j+r(g)-(r(g_2)+r(g_1)) = j+r(g)-\alpha-n_2+n_1 = k'-n_2$. Denote $l = j+r(g_4)$ and let $k' \in U$ mean that k' is divisible by n_3. With this notation we have $L_{j',g} \subseteq \sum_{i=1}^{2t\alpha+t}\sum_{k',l:k'\in U,l+r(g_3)=k'-n_2}((A(j'\cdot n_3+\beta+i)\otimes A(l))\xi(g_3)(A\otimes A)$.

Recall that $2n_1 < c < \beta \le n_2 - 10tn_1$ and $\alpha < 2n_1$, $g_3 \subseteq \sum_{i'=1}^{2tn_1+t} A(i')\otimes A(n_1)$, $\alpha \le 2n_1$. Hence for $i \le 2t\alpha+t, i' \le 2tn_1+2t$ the interval $[\beta+i, \beta+i+i']$ is disjoint with any interval $[kn_2-n_1, kn_2+n_1]$. Recall that and that n_3 is divisible by n_2, so interval $[j'n_3+\beta+i, j'n_3+\beta+i+i']$ is disjoint with any interval $[kn_2-n_1, kn_2+n_1]$.

Therefore, $L_{j',g'} \subseteq Q'(\xi(T'(f,n_1)),n_1,n_2) = Q_3$, as required. □

We are now ready to prove Theorem 3.1.

Proof of Theorem 3.1. If K is a countable field then we can ennumerate elements of $A\otimes A$, as f_1, f_2, in a such way that: all elements of $A\otimes A$ are listed; some f_i can be zero; $f_i \subseteq \sum_{i,j\le i-1} A(i)\otimes A(j)$ for all $i \ge 2$, and moreover $f_1 = f_2 = 0$. Let N be the ideal generated by $f_i^{20ia_{i+1}}$ and $\xi(f_i^{20ia_{i+1}})$ for all $i \ge 2$. By Theorem 12.1, applied for $f = f_n$, $n_1 = a_{n-1}$, $n_2 = a_n$, $n_3 = a_{n+1}$, $t = n-1$, the two sided ideal generated by $f_n^{100a_{n+2}}$ and $\xi(f_n^{100a_{n+2}})$ is contained in $M_{n+1} + Q_n + Q_{n-1} + Q'_{n-1}$, where linear spaces F_n, F'_n and G_n are defined as follows. With notation as in Theorem 12.1, for each $n \ge 1$ denote

$$F_n = F(f_n,a_{n-1},a_n,a_{n+1})+F'(f_n,a_{n-1},a_n,a_{n+1})+T(f_{n+1},a_n)+\xi(T(f_{n+1},a_n)),$$

$$F'_n = T'(f_{n+1},a_n)+\xi(T'(f_{n+1},a_n)), G_n = G(f_n,a_{n-1},a_n).$$

We will show that F_n, F'_n, G_n satisfy assumptions of our theorem. By Theorem 12.1, $G_n \subseteq A(a_n)\otimes A(a_n)$ and $\dim_K G_n < a_n^3 2^{500na_{n-1}} < a_{n+2}^{100n}2^{500na_{n-1}}$. Observe that by Lemmas 11.3 and 11.4, we have $\dim_K F(f_n,a_{n-1},a_n,a_{n+1})$, $\dim_K F'(f_n,a_{n-1},a_n,a_{n+1}) \le 2^{40n\cdot a_{n-1}}a_n$ and By Lemmas 9.3, 9.5 and 9.6, we have $\dim_K T(f_{n+1},a_n)$, $\dim_K T'(f_{n+1},a_n) \le max(2^{4n+3}a_n^3n^3, 2^{2n^2+1})$ where $max(a,b)$ denotes the maximum of a and b. Therefore $\dim_K F_n, \dim_K F'_n \le a_{n+2}^{100n}2^{500n\cdot a_{n-1}}$. By Lemmas 11.3 and 11.4, and since $a_n > 100na_{n-1}$ for every i, we have $F(f_n,a_{n-1},a_n,a_{n+1}), F'(f_n,a_{n-1},a_n,a_{n+1}) \subseteq A(a_n)\otimes\sum_{j>\frac{a_n}{10n}}^{10na_n} A(j)$. By Lemmas 9.3, 9.5 and 9.6, we have $T(f_{n+1},a_n) \subseteq A(a_n)\otimes\sum_{i\ge\frac{a_n-n}{2n}}^{2na_n+n} A(i)$

and $T'(f_{n+1}, a_n) \subseteq \sum_{i \geq \frac{a_n - n}{2n}}^{2na_n+n} A(i) \otimes A(a_n)$. It follows that $F_n \subseteq A(a_n) \otimes \sum_{j > \frac{a_n}{10n}}^{10na_n} A(j)$ and $F'_n \subseteq \sum_{j > \frac{a_n}{10n}}^{10na_n} A(j) \otimes A(n)$, since $n < \frac{a_n}{2}$. Notice that we can put $G_1 = 0$, $G_2 = 0$, $F_1 = 0$ because $f_1, f_2 = 0$.

Assume now that K is an uncountable field, as A has two generators , it follows that set $M \otimes M$ is countable, where $M \subseteq A$ is a set of monomials in A. Therefore, the set of finite subsets of $M \otimes M$ is also countable. If H is a subset of $M \otimes M$ by KH we will mean the set of all linear combinations of elements from H.

Let $H_2, H_3, \ldots$ be finite subsets of $M \otimes M$ such that for each $i \geq 2$, the cardinality of H_i is less than i and $H_i \subseteq \sum_{i,j \leq i-1} A(i) \otimes A(j)$ and for all $i \geq 2$. Moreover some sets H_i may be zero. Observe that we can list all finite subsets of $M \otimes M$ as sets $H_2, H_3, \ldots$, because we allow some sets H_i to be zero. Moreover, we can assume that $H_1 = H_2 = H_3 = \emptyset$.

For a given set H and a natural number n let $P_n(H) = \{f^n : f \in H\}$. Then $K \cdot P_n(H)$ is the linear space spanned by elements from $P_n(H)$.

We now claim that for every $n \geq 2$ there is a set $\bar{H}_n \subseteq KH_n$ and such that the cardinality of $\bar{H}_n$ is less than $(20na_{n+1})^n$ and moreover $K \cdot P_{20na_{n+1}}(\bar{H}_n) = K \cdot P_{20na_{n+1}}(KH_n)$. To see that let $f \in K \cdot H_n$, then $f = \sum_{i=1}^{n} \alpha_i m_i$ where $\alpha_i \in K$, $m_i \in H$. Let $w(j_1, j_2, \ldots, j_n)$ be the sum of all products of length $20na_{n+1}$ of elements from the set H which are products of exactly j_i elements m_i for every $i \leq n$. Then $f^{20na_{n+1}}$ is a linear combination of elements $w(j_1, j_2, \ldots, j_n)$ for various $j_1, j_2, \ldots, j_n$, with $j_1 + \ldots + j_n = 20na_{n+1}$. Because there are less than $(20na_{n+1})^n$ such choices of elements $i_1, i_2, \ldots, i_n$, it follows that the dimension of the linear space $P_{20na_{n+1}}(KH_n)$ is less than $(20na_{n+1})^n$. Therefore, there is a subset $\bar{H}_n \subseteq KH_n$ with cardinality less than $(20na_{n+1})^n$ and such that $K \cdot P_{20na_{n+1}}(\bar{H}_n) = K \cdot P_{20na_{n+1}}(KH_n)$, as required. Notice than then also $K \cdot \xi(P_{20na_{n+1}}(\bar{H}_n)) = K \cdot \xi(P_{20na_{n+1}}(KH_n)$,

We will now proceed in a similar way as in the case of countable fields at the beginning of this proof. We can assume that $H_1 = H_2 = H_3 = \emptyset$ and put $F_1 = 0$, $G_1 = 0$, $G_2 = 0$. Observe that by Theorem 12.1 applied for $f \in \bar{H}_n$, $n_1 = a_{n-1}$, $n_2 = a_n$, $n_3 = a_{n+1}$, $t = n - 1$, the two sided ideal generated by elements from the set $\{f^{100a_{n+2}} : f \in \bar{H}_n\}$ and from the set $\{\xi(f)^{100a_{n+2}} : f \in \bar{H}_n\}$ is contained in $M_{n+1} + Q_n + Q_{n-1} + Q'_{n-1}$, where linear spaces F_n, F'_n and G_n are defined as follows. $F_1 = 0$, $G_1 = 0$, $G_2 = 0$, and $F_n = \{F(f, a_{n-1}, a_n, a_{n+1}) + F'(f, a_{n-1}, a_n, a_{n+1}) + T(g, a_n) + \xi(T(g, a_n)) : f \in \bar{H}_n, g \in \bar{H}_{n+1}\}$. Moreover,

$$F'_n = \{T'(g, a_n) + \xi(T'(g, a_n) : g \in \bar{H}_{n+1}, \},$$

$$G_n = \{G(f, a_{n-1}, a_n) : f \in \bar{H}_n\}.$$

We will show that F_n, F'_n, G_n satisfy the assumptions of our theorem. By Theorem 12.1, $G_n \subseteq A(a_n) \otimes A(a_n)$ and for every $g \in \bar{H}_{n+1}$, $\dim_K G(g, a_{n-1}, a_n) < a_n^3 2^{500na_{n-1}}$. Hence, $\dim_K G_n < (a_n^3 2^{500na_{n-1}}) \cdot (20(n+1)a_{n+2}^{n+1}) < a_{n+2}^{100n} 2^{500na_{n-1}}$ (because $\dim_K \bar{H}_{n+1} \leq 20(n+1)a_{n+2}^{n+1}$). By Lemmas 11.3, 9.6 and 11.4, we have for each $f \in \bar{H}_n$, $\dim_K F(f, a_{n-1}, a_n, a_{n+1}), \dim_K F'(f, a_{n-1}, a_n, a_{n+1}) \leq 2^{40n \cdot a_{n-1}} a_n$ and By Lemmas 9.3, 9.5 and 9.6, we have for every $g \in \bar{H}_{n+1}$, $\dim_K T(g, a_n), \dim_K T'(g, a_n) \leq max(2^{4n+3} a_n^3 n^3, 2^{2n^2+1})$. Recall that $\dim_K \bar{H}_n$, $\dim_K \bar{H}_{n+1} \leq 20(n+1)a_{n+2}^{n+1}$, it follows that $\dim_K F_n, \dim_K F'_n \leq a_{n+2}^{100n} 2^{500n \cdot a_{n-1}}$. Similarly as in the countable field case, $F_n \subseteq A(a_n) \otimes \sum_{j > \frac{a_n}{10n}}^{10na_n} A(j)$ and $F'_n \subseteq \sum_{j > \frac{a_n}{10n}}^{10na_n} A(j) \otimes A(n)$. Therefore, sets F_n, F'_n, G_n satisfy thesis of our theorem.

Let N be the ideal generated in $A \otimes A$ by sets $P_{20na_{n+1}}(\bar{H}_n)$, for $n = 2, 3,$. By the above reasoning, $N \subseteq \sum_{i=1}^{\infty} M_i + Q_i$. Moreover, if $f \in A \otimes A$ then $A \in H_i$ for some i. Then $f^{20ia_{i+1}} \in P_{20ia_{i+1}}(KH_i) \subseteq K \cdot P_{20ia_{i+1}}(\bar{H}_i) \subseteq N$. Similarly, $\xi(f^{20ia_{i+1}}) \in \xi(P_{20ia_{i+1}}(KH_i)) \subseteq K \cdot \xi(P_{20ia_{i+1}}(\bar{H}_i)) \subseteq N$.

Acknowledgements This research was supported by ERC Advanced grant Coimbra 320974. The author would like to thank Adam Chapman, Natalia Iyudu, Jerzy Matczuk, Uzi Vishne and Michał Ziembowski for their comments on the paper's introduction. The author is also grateful to Pace Nielsen, as a simple exposition from his paper [14] was used by her to simplify the proof of Theorem 3.1. The author is especially grateful to Andre Leroy and the organisers of "Nord Pas de Calais/Belgium congress of Mathematics" for their hospitality during her stay in Valenciennes, Mons and Lens, in October and November 2013 where the first part of this paper was prepared.

References

[1] B. Amberg, L. Kazarin, *On the adjoint group of a finite nilpotent p-algebra,* Journal of Mathematical Sciences November 25, 2000, Volume 102, Issue 3, pp 3979-3997.

[2] G. M. Bergman,Radicals, tensor products, and algebraicity, Israel Mathematical Conference Proceedings1 (1989), 150-192.

[3] L. Bokut, Yu. Chen, Yo. Chen, *Composition-Diamond lemma for tensor product of free algebras,* Journal of Algebra, Volume 323, Issue 9, 1 May 2010, Pages 2520-2537.

[4] M. A. Chebotar, P. H. Lee, E. R. Puczylowski, *A linear algebra approach to Koethe's problem and related questions,* Linear and Multilinear Algebra 2013, Volume 61, Issue 5, 635-645.

[5] M. Ferrero, E.R. Puczylowski, *Prime ideals and radicals of centered extensions and tensor product,* Israel Journal of Mathematics 1996, Volume 94, Issue 1, pp 381-401.

[6] E. S. Golod, Some problems of Burnside type, Proc. I.C.M., Moscow, 1966, 284-289.

[7] E. S. Golod and I. R. Shafarevich, On class field towers, Izv. Akad. Nauk SSSR Ser. Mat. 28 (1964), 261272; English transl., Amer. Math. Soc. Transl. (2) 48 (1965), 91-102.

[8] E. Jespers and E. R. Puczyłowski, *On ideals of tensor products,* Journal of Algebra140 (1991), 124-130.

[9] J. Krempa, *On semisimplicity of tensor products,* Proc. 1978 Antwerp Conference, Lecture Notes in Pure and Applied Mathematics, Vol. 51, Dekker, New York, 1979, pp. 105-122.

[10] T.Y. Lam, A first course in Noncommutative Ring Theory, Springer Graduate Series 2002.

[11] John Lawrence, *The Jacobson radical of tensor products,* Quarterly Journal of Mathematics (1991) 42 (1), 203-208.

[12] John Lawrence, *When is the tensor product of algebras local?* Proc. Amer. Math. Soc. 58 (1976), 22-24.

[13] P. Kanwar, A. Leroy, J. Matczuk, *Clean elements in polynomial rings,* preprint.

[14] Pace P. Nielsen, *Simplifying Smoktunowicz's extraordinary example*, Communications in Algebra, Volume 41, Issue 11, 2013, 4339-4350.

[15] W. K. Nicholson and J. F. Watters, *On tensor products and extended centroids,* Proceedings of the American Mathematical Society 88 (1983), 215-217.

[16] E.R. Puczylowski, *Some questions concerning radicals of associative rings,* in Theory of Radicals. Colloq. Math. Soc. Janos Bolyai, 61 (1993), pp. 209-227.

[17] E. R. Puczyłowski,*Nil algebras with nonradical tensor square,* Proceedings of the American Mathematical Society 103 (1988), 401-403.

[18] A.Regev, A.Regev, *The Golod-Shafarevich theorem without Hilbert series*, Contemporary Mathematics, 2009, Proceedings of the International Conference : Groups, Rings and Group Rings, edited by A. Giambruno, Csar Polcino Milies, Sudarshan K. Sehgal 257-264.

[19] J. C. Robson, *Prime ideals in intermediate extensions,* Proceedings of the London Mathematical Society 44 (1982), 372-384.

[20] R. Resco, L. W. Small and A. Wadsworth, *Tensor products of division rings and finite generation of subfields,* Proc.Amer.Math.Soc. 77 (1979) 7-10.

[21] Agata Smoktunowicz, *Polynomial rings over nil rings need not be nil*, J. Algebra Volume 233, Issue 2, 2000, 427-436.

[22] A. Smoktunowicz, L. Bartholdi, *Images of Golod-Shafarevich algebras with small growth,* Quarterly Journal of Mathematics, (2013) doi: 10.1093/qmath/hat005.

[23] M. Tousi, S. Yassemi, *Tensor products of clean rings,* Glasgow Mathematical Journal, Volume 47, Issue 03, 2005, 501-503.

[24] P. Vamos, *On the minimal prime ideals of a tensor product of two fields,* Math. Proc. Cambridge Philos. Soc. 84 (1978), 25-35.